

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

C12N 15/12, C07K 14/78, C12Q 1/68,
C12N 15/62, C07K 16/18, A61K 38/39

A1

(11) International Publication Number:

WO 99/29855

(43) International Publication Date:

17 June 1999 (17.06.99)

(21) International Application Number:

PCT/US98/26057

(22) International Filing Date:

8 December 1998 (08.12.98)

(30) Priority Data:

60/067,888	8 December 1997 (08.12.97)	US
60/082,663	22 April 1998 (22.04.98)	US
60/108,536	16 November 1998 (16.11.98)	US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications

US	60/067,888 (CIP)
Filed on	8 December 1997 (08.12.97)
US	60/082,663 (CIP)
Filed on	22 April 1998 (22.04.98)
US	60/108,536 (CIP)
Filed on	16 November 1998 (16.11.98)

(71) Applicant (for all designated States except US): BETH ISRAEL
DEACONESS MEDICAL CENTER [US/US]; 330 Brook-
line Avenue, Boston, MA 02215 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): SUKHATME, Vikas, P.
[US/US]; 36 Sky View Circle, Newton Center, MA 02159
(US).(74) Agents: HOGLE, Doreen, M. et al.; Hamilton, Brook, Smith &
Reynolds, P.C., Two Militia Drive, Lexington, MA 02421
(US).(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR,
BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP,
KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK,
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU,
ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG,
ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI,
FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE,
SN, TD, TG).

Published

*With international search report.
Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: MUTANTS OF ENDOSTATIN, "EM 1" HAVING ANTI-ANGIOGENIC ACTIVITY AND METHODS OF USE THEREOF

(57) Abstract

Described herein are novel mutants of endostatin, one of which, designated "EM 1", has anti-angiogenic activity similar or superior to that of wild type endostatin. The invention relates to the discovery of an isolated anti-angiogenic peptide, wherein the C-terminal end of the peptide comprises the amino acid sequence SYIVLCIE, which has anti-angiogenic properties. Designated "EM 1", this protein comprises a mutated endostatin protein, where the mutation comprises a deletion of nine consecutive amino acids from the C-terminus of the mutated endostatin protein (e.g., NSFMTSFSK). EM 1 terminates in the amino acid sequence SYIVLCIE. The invention also comprises isolated polynucleotides encoding EM 1, operably linked to expression sequence, and host cells transformed with such a construct. Antibodies to EM 1 are also disclosed. The invention also relates to processes for producing EM 1, fusion proteins containing EM 1, and compositions comprising EM 1 or fusion products thereof. The invention also discloses methods of producing polypeptides encoding EM 1.

AN

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MUTANTS OF ENDOSTATIN, "EM 1" HAVING ANTI-ANGIOGENIC ACTIVITY AND METHODS OF USE THEREOF

RELATED APPLICATIONS

This application claims priority to application 60/067,888, filed December 8, 1997, 60/082,663, filed April 22, 1998, and 60/108,536, filed November 16, 1998, the entire teachings of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

The prognosis for metastatic cancer remains highly unfavorable. Despite advances in radiation therapy and chemotherapy, the long term survival of treated patients has shown only marginal improvement over the past few decades. The lack of significant treatment options available for metastatic cancers emphasizes the need to focus on the development of novel therapeutic strategies. In this regard, targeting tumor vasculature of solid tumors has recently shown promising results in several animal model systems (Baillie *et al.* (1995) *Br. J. Cancer* 72:257-67; Bicknell, R. (1994) *Ann. Oncol.* 5 (Suppl.) 4:45-50; Fan *et al.* (1995) *Trends Pharmacol. Sci.* 16:57-66; Thorpe, P.E. and Burrows, F.J. (1995) *Breast Cancer Res. Treat.* 36:237-51; Burrows, F.J. and Thorpe, P.E. (1994) *Pharmacol. Ther.* 64:155-74). In a nude mouse model, for instance, introduction of a wild type VHL gene into 786-0 cells, a RCC tumor cell line, inhibited tumor growth (Iliopoulos *et al.* (1995) *Nat. Med.* 1:822-26) and angiogenesis.

The growth of solid tumors beyond a few mm³ depends on the formation of new blood vessels (Folkman, J. (1971) *N. Engl. J. Med.* 285:1182-86). Numerous studies have shown that both primary tumor and metastatic growth are angiogenesis-dependent (Folkman, J. (1971) *N. Engl. J. Med.* 285:1182-86; Folkman, J. (1972) *Ann. Surg.* 175:409-16; Folkman, J. and Shing, Y. (1992) *J. Biol. Chem.* 267:10931-34; Folkman, J. (1996) *Sci. Am.* 275:150-54). A number of angiogenesis inhibitors have been identified. Certain ones, such as platelet factor-4 (Maione *et al.* (1990) *Science* 247:77-79; Gupta *et al.* (1995) *Proc. Natl. Acad. Sci.*

(USA) 92:7799-7803), interferon α , interferon-inducible protein-10, and PEX (Angiolillo *et al.* (1995) *J. Exp. Med.* 182:155-62; Strieter *et al.* (1995) *Biochem. Biophys. Res. Commun.* 210:51-57; Brooks *et al.* (1998) *Cell* 92:391-400), are not "associated with tumors," whereas two others, angiostatin and endostatin, are

5 "tumor-associated" (O'Reilly *et al.* (1994) *Cell* 79:315-28; O'Reilly *et al.* (1997) *Cell* 88:277-85). Angiostatin, a potent endogenous inhibitor of angiogenesis generated by tumor-infiltrating macrophages that upregulate matrix metalloelastase (Dong *et al.* (1997) *Cell* 88:801-10), inhibits the growth of a wide variety of primary and metastatic tumors (Lannutti *et al.* (1997) *Cancer Res.* 57:5277-80; O'Reilly *et al.*

10 (1994) *Cold Spring Harb. Symp. Quant. Biol.* 59:471-82; O'Reilly, M.S., (1997) *Exs.* 79:273-94; Sim *et al.* (1997) *Cancer Res.* 57:1329-34; Wu *et al.* (1997) *Biochem. Biophys. Res. Commun.* 236:651-54).

Recently, O'Reilly, *et al.* ((1997) *Cell* 88:277-85) isolated endostatin, an angiogenesis inhibitor from a murine hemangioendothelioma cell line (EOMA).

15 Circulating levels of a fragment of human endostatin have been detected in patients with chronic renal insufficiency with no detectable tumor, but this fragment had deletions, and no anti-angiogenic activity (Standker *et al.* (1997) *FEBS Lett.* 420:129-33). The amino terminal sequence of endostatin corresponds to the carboxy terminal portion of collagen XVIII. Endostatin is a specific inhibitor of endothelial

20 proliferation and angiogenesis. Systemic administration of non-refolded precipitated protein expressed in *Escherichia coli* caused growth regression of Lewis lung carcinoma, T241 fibrosarcoma, B16 melanoma and EOMA (O'Reilly *et al.* (1997) *Cell* 88:277-85) cells in a xenograft model. Moreover, no drug resistance was noted in three of the tumor types studied. Repeated cycles of administration with

25 endostatin have been reported to result in tumor dormancy (Boehm *et al.* (1997) *Nature* 390:404-407).

The results from these studies open new avenues for treatment of cancer and provide promising routes for overcoming the drug resistance often seen during chemotherapy. However, in all of these investigations, a non-refolded precipitated

30 form of the inhibitor protein was administered in the form of a suspension to tumor

bearing animals. In addition, large amounts of protein were required to cause tumor regression and to lead to tumor dormancy. As pointed out by Kerbel ((1997) *Nature* 390:335-36), oral drug equivalents of these proteins are needed. Mechanistic investigations could be undertaken if recombinant forms of these proteins were
5 available in soluble form. Moreover, initial testing could be done *in vitro* with soluble protein before studying its efficacy under *in vivo* conditions. Furthermore, there have been reports that despite the great promise held by these proteins, evaluation of their clinical potential is stymied due to difficulties in producing enough of the protein to test, and inconsistent test results regarding their anti-
10 angiogenic properties (King, R.T. (1998) *Wall Street J.*, page 1 Nov. 12; Leff, D.N. (1998) *BioWorld Today* 9:1, Oct. 20). There clearly exists at the present time a great need for a method of producing soluble forms of anti-angiogenic proteins in large amounts, and which have reliable properties *in vitro* and *in vivo*.

SUMMARY OF THE INVENTION

15 Described herein are novel mutants of endostatin, one of which, designated "EM 1," has anti-angiogenic activity similar or superior to that of wild type endostatin.

The invention relates to the discovery of an isolated anti-angiogenic peptide, wherein the C-terminal end of the peptide comprises the amino acid sequence
20 SYIVLCIE, which has anti-angiogenic properties. Designated "EM 1," this protein comprises a mutated endostatin protein, where the mutation comprises a deletion of nine consecutive amino acids from the C-terminus of the mutated endostatin protein (*e.g.*, NSFMTSFSK). EM 1 terminates in the amino acid sequence SYIVLCIE. The invention also comprises isolated polynucleotides encoding EM 1, operably linked to
25 expression sequences, and host cells transformed with such a construct. Antibodies to EM 1 are also disclosed.

The invention also relates to processes for producing EM 1, fusion proteins containing EM 1, and compositions comprising EM 1 or fusion products thereof. The invention also discloses methods of producing polypeptides encoding EM 1.

In addition, the invention comprises methods for inhibiting angiogenic activity in mammalian tissue, comprising contacting the tissue with a composition comprising the EM 1, particularly to inhibit angiogenesis, which occurs in many diseases and conditions, including cancer.

5 The invention also discloses use of EM 1 to induce apoptosis, or antibodies of EM 1 to prevent apoptosis. The invention further discloses use of EM 1 in methods of gene therapy. The cells targeted may be any mammalian cells, particularly lymphocytes, blood cells, TIL cells, bone marrow cells, vascular cells, tumor cells, liver cells, muscle cells, and fibroblast cells.

10 BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a diagram of the endostatin nucleotide sequence (SEQ ID NO:1). The polynucleotide encoding EM 1 comprises the polynucleotide sequence through nucleotide 525. The polynucleotide encoding EM 2 comprises the polynucleotide sequence through nucleotide 501.

15 Fig. 2 is a diagram of the translation (SEQ ID NO:2) of the nucleic acid sequence of Fig. 1. EM 1 comprises the amino acid sequence through amino acid 175. EM 2 comprises the amino acid sequence through amino acid 167.

Fig. 3 is a graph showing the results of elution from an Ni-NTA column. The fraction number is shown along the x-axis, and the absorbance at 280 nm for each fraction (O) is on the left y-axis. For each fraction, the pH of the eluting buffer (■) is shown on the right y-axis.

Fig. 4 shows a 12 % non-reducing SDS-PAGE gel of protein produced from the prokaryotic expression system. Sizes in kDa are shown on the left, and the first lane contains size markers. Lane 2 contains crude protein, lanes 3 and 4 contain samples from fractions 7 and 8, which were eluted at pH 6.3. Lanes 5 and 6 contain samples from fractions 21 and 22, eluted at pH 4.0. Lane 7 contains a sample from fraction 22 reduced with DTT.

Fig. 5 is a graph showing the purification of soluble mouse endostatin expressed in yeast using a heparin-agarose column. The fraction number is shown along the x-axis, and the absorbance at 280 nm for each fraction (O) is on the left y-

axis, and the concentration of NaCl used to elute each fraction (■) is shown on the right y-axis.

Fig. 6 shows a 12% non-reducing SDS-PAGE gel of purified recombinant soluble mouse endostatin from a heparin-agarose column. Sizes in kDa are shown on the left, and the first lane contains size markers. Lane 2 contains crude protein, lane 3 contains unbound protein, lane 4 contains wash, and lanes 5 and 6 contain samples from fractions 9 and 10, respectively, which were both eluted with 0.3ml NaCl. Lanes 7 and 8 contain samples from fractions 21 and 22, respectively, which were both eluted with 0.6 M NaCl.

Fig. 7 is a graph showing the elution profile of soluble His.endostatin expressed in yeast using a Ni-NTA column. The fraction number is shown along the x-axis, and the absorbance at 280 nm for each fraction (○) is on the left y-axis, and the concentration of imidazole (mM) used to elute each fraction (■) is shown on the right y-axis.

Fig. 8 shows a 12% non-reducing SDS-PAGE gel of selected fractions of soluble His.endostatin expressed in yeast. Sizes in kDa are shown on the left, and the first lane contains size markers. Lane 2 contains crude protein, lane 3 contains flowthrough (*i.e.*, unbound protein), and lane 4 contains wash. Lane 5 contains a sample of fraction 9, which was eluted at 10 mM imidazole. Lanes 6 and 7 contain samples from fractions 35 and 36, respectively, which were eluted with 50 mM imidazole, and lanes 8 and 9 contain samples of fractions 53 and 54, respectively, which were eluted with 100 mM imidazole.

Fig. 9 shows a Western blot analysis of recombinant mouse endostatin expressed from bacteria and yeast. Lane 1 contains bacterially-expressed His.endostatin, lane 2 contains endostatin expressed in yeast, and lane 3 contains yeast-produced His.endostatin.

Fig. 10 is a graph depicting the results of an endothelial cell proliferation assay. The purified mouse endostatin expressed from yeast was tested for its ability to inhibit (methyl- ^3H) thymidine incorporation in C-PAE cells. The concentration of endostatin (100 ng to 10000 ng) is shown on the x-axis, and the incorporation of

^3H -thymidine is shown on the y-axis. Incorporation for yeast-derived soluble endostatin (○) and yeast-derived soluble His.endostatin (■) dropped steadily with increasing concentration of endostatin.

Fig. 11 is a bar chart showing the effects of recombinant mouse endostatin on non-endothelial cells. Open bars refer to 786-0 cells, and shaded bars refer to A₄₉₈ cells. Both are renal carcinoma cell lines, stimulated with bFGF (3 ng/ml) in 2% serum.

Figs. 12A and 12B are a pair of photographs showing inhibition of endothelial cell (ECV304) migration by soluble mouse endostatin using bFGF (25 ng/ml) as a stimulus. Fig. 12A shows migrated endothelial cells in the control (+ bFGF, no endostatin), and Fig. 12B shows migrated endothelial cells treated with endostatin (20 $\mu\text{g/ml}$) with BFGF.

Fig. 13 is a bar chart showing inhibition of endothelial cell migration with different concentrations of endostatin. Relative cell migration is shown on the y-axis, and treatment (control, 25 ng/ml bFGF, and endostatin at 20, 10, 5, 2.5, and 1 $\mu\text{g/ml}$) on the x-axis.

Figs. 14A, 14 B, and 14C are photomicrographs showing the inhibition of angiogenic response mediated by VEGF (250 ng/pellet) in the presence of endostatin. Fig. 14A is the negative control, Fig. 14B is the positive control (VEGF), and Fig. 14C shows the effect of endostatin plus VEGF.

Figs. 15A and 15B are a pair of bar charts showing the inhibition of VEGF (top panel) and bFGF (bottom panel) mediated angiogenic response by endostatin (20, 10, 5, 1, and 0 $\mu\text{g/mesh}$) in the CAM assay. Both charts show a steady increase of inhibition of angiogenesis in response to increasing concentrations of endostatin.

Fig. 16 is a bar chart showing neutralization of the inhibitory effect of mouse endostatin by polyclonal antiserum in the endothelial proliferation assay.

Incorporation of ^3H -thymidine is shown on the y-axis, and treatment (control, 10 μg endostatin, 10 μg endostatin + antiserum, 5 μg endostatin, 5 μg endostatin + antiserum, pre-immune serum, endostatin antiserum, and endostatin IgG) on the x-

axis).

Figs. 17A and 17B are a pair of photographs showing the results of a CAM assay, demonstrating neutralization of endostatin inhibitory activity by polyclonal antiserum. Fig. 17A shows the effect of VEGF and endostatin (10 $\mu\text{g}/\text{pellet}$), and Fig. 17B shows the effect of endostatin (10 $\mu\text{g}/\text{pellet}$) plus polyclonal antiserum plus VEGF.

Fig. 18 is a graph showing the inhibition of 786-0 tumor growth by systemic treatment with recombinant endostatin. Time in days after treatment is shown on the x-axis and tumor volume in mm^3 is shown on the y-axis. Intraperitoneal injection of endostatin was given at 10 mg/kg/day, starting on day 1 (arrow). Each time point represents the average of five mice in each group and the error bar represents S.E.M. Treatments are control PBS (O), endostatin from yeast (\bullet), His.endostatin from yeast (x) and His.endostatin from bacteria (\square).

Figs. 19A through 19E are a set of photographs of 786-0 tumors treated with recombinant endostatin. At the end of the treatment period, tumors from control and treated groups were examined grossly under a dissecting microscope. Figs. 19A and 19B are control tumors, Fig. 19C shows a tumor treated with yeast-derived endostatin, Fig. 19D shows the effect of His.endostatin from bacteria, and Fig. 19E shows a tumor treated with His.endostatin from yeast.

Fig. 20 is a graph showing the effects of endostatin mutants on athymic nude mice 786-0 tumors. Days after treatment is shown on the x-axis, and tumor volume on the y-axis. Each time point represents the average of five mice in each group. Treatments were control PBS (O), wild type His.endostatin from bacteria (dotted line, \square), EM 1 from bacteria (Δ), and EM 2 from bacteria (solid line, \blacklozenge). EM 1 and EM 2 both contain N-terminus His.tags. Intraperitoneal injection was started on day 1 (arrow).

Fig. 21 is a bar graph showing increased caspase 3 activity due to endostatin treatment. Absorbance at 405 nm is shown on the y-axis, and treatments (control, $\text{TNF-}\alpha$ (10 ng/ml), endostatin (10 $\mu\text{g}/\text{ml}$) are shown on the x-axis. The pairs of bars for each treatment the A_{405} reading in the presence (open bars) or absence (shaded bars) of the inhibitor DEVD-fmk.

Fig. 22 is a bar graph showing caspase 3 activity in non-endothelial cells. Absorbance at 405 nm is shown on the y-axis, and x-axis displays treatments (control, control + DEVD, endostatin (10 μ g/ml), endostatin (10 μ g/ml) + DEVD) for NIH3T3 and H9c2 (2-1)-myoblast cells, respectively.

5 Fig. 23 is a bar graph showing quantitative determination of apoptosis, as determined by the TUNEL assay. The treatments are shown on the x-axis, and are endostatin-treated adherent cells, endostatin-treated suspension cells, control adherent cells, and TNF- α -treated adherent cells. Percentage of apoptotic cells are shown on the y-axis.

10 Figs. 24A and 24B are a Western blot analyses of C-PAE cell lysate for Bcl-2 protein levels, and an immunoblot detecting total cell lysate for Bax expression levels, respectively. C-PAE cells were treated with either no endostatin (-) or endostatin (10 μ g/ml) (+) for the indicated period of time 0, 12, 24, 28 hours). Actin probing is also shown.

15 Figs 25A, 25B, 25C and 25D are a set of two Western blot analyses (Figs. 25A and 25B) and two immunoblots (Figs. 25C and D) of non-endothelial cell lysate for Bax protein levels. Cells were treated with either no endostatin (-) or endostatin (10 μ g/ml) (+) for the indicated period of time 0, 12, 24, 28, 32 hours). Actin probing is also shown. Fig. 25A: NIH3T3 cell lysate; Fig. 25B: IMR-90 cell
20 lysate; Fig. 25C: C-PAE cell lysate; Fig. 25D: NIH3T3 cell lysate.

Figure 26A-B is a chart showing the constructs, primers, cloning sites, and vectors, used to clone and express various anti-angiogenic proteins. The amino acid sequences of the expressed proteins are also given.

DETAILED DESCRIPTION OF THE INVENTION

25 A wide variety of diseases are the result of undesirable angiogenesis. Put another way, many diseases and undesirable conditions could be prevented or alleviated if it were possible to stop the growth and extension of capillary blood vessels under some conditions, at certain times, or in particular tissues. Several anti-angiogenic proteins have been discovered (e.g., angiostatin, endostatin), problems

have been reported regarding (1) the ability to produce the proteins in sufficient quantity to allow for proper testing of their properties, and (2) the reproducibility of the anti-angiogenic properties attributed to these proteins.

The present invention encompasses endostatin mutants, referred to herein as
5 EMs. Specifically encompassed are two mutants of endostatin, designated "EM 1"
and "EM 2". These mutants were tested against whole endostatin. The mutants
showed very different activity. Unexpectedly, one mutant ("EM 1") performed as
well or better than whole endostatin, and the other ("EM 2") showed loss of anti-
angiogenic activity. In a nude mouse model, growth of renal cell cancer (RCC) was
10 suppressed by systemic administration of EM 1 at a rate of 20 mg/kg body weight.
The inhibition of tumor growth is comparable to the inhibition obtained with wild-
type endostatin. The difference in activity between EM 1 and EM 2 is surprising,
given that there is a difference between them of only eight amino acid residues.
EM 1 provides an advantage in treatment of angiogenic diseases in that increasingly
15 smaller peptides are more potent on a weight basis, and may be able to better
penetrate tissues.

In the present invention, EM 1, a novel anti-angiogenic protein, and a
deletion mutant of endostatin, is described, as well as fragments, derivatives, fusion
proteins and antibodies thereof. Methods of making the above are also described.
20 Also disclosed are therapeutic compositions comprising EM 1, and methods for
using those compositions. Polynucleotides encoding EM 1 are also described, as well
as vectors and host cells comprising those polynucleotides. Compositions
containing EM 1 as a biologically active ingredient are also described, as well as
methods for using EM 1 to inhibit angiogenic activity in mammalian tissues, such as
25 in treating diseases and conditions characterized by angiogenesis. The present
invention includes compositions and methods for the detection and treatment of
diseases and conditions that are mediated by or associated with angiogenesis. In
addition, the invention includes use of EM 1 to induce apoptosis in a cell or tissue,
and antibodies to EM 1 to inhibit apoptosis in a cell or tissue.

Specifically, EM 1 is a deletion mutant of endostatin, where the last nine amino acid residues have been deleted. EM 1 exists naturally as part of the collagen Type XVII molecule, but it can be produced recombinantly, *e.g.*, the polynucleotide sequence (Fig. 1, SEQ ID NO:1) encoding EM 1 protein (Fig. 2, SEQ ID NO:2) can
 5 amplified, *e.g.*, with the forward and reverse primers listed in Table 1, below. The template nucleic acid used for the amplification can be from any mammal. Also encompassed by the present invention is mammalian EM1, fragments, mutants, derivatives or fusion proteins thereof.

Table 1. Constructs and primer sequences used to amplify anti-angiogenic proteins.

10	Construct Name	Primer Sequence
	pET17bhis.mendo	5'-GGC ATA TGC ATA CTC ATC AGG ACT TT-3' (up) (SEQ ID NO.:)
		5' AAC TCG AGC TAT TTG GAG AAA GAG GT-3' (down) (SEQ ID NO.:)
	pET28a/mendo	5'-GGC ATA TGC ATA CTC ATC AGG ACT TT-3' (up)
		5'-AAG CGG CCG CCT ATT TGG AGA AAG AGG T-3' (down) (SEQ ID NO:)
	pET28a/EM-1	5' TTC CAT ATG CAT ACT CAT CAG GAC TTT CAG CCA-3' (up) (SEQ ID NO:)
		5' TTA GCG GCC GCC TAC TCA ATG CAC AGG ACG ATG TA-3' (down) (SEQ ID NO:)
	pET28a/EM-2	5' TTC CAT ATG CAT ACT CAT CAG GAC TTT CAG CCA-3' (up) (SEQ ID NO:)
		5' TTA GCG GCC GCC TAG TTG TGG CAG CTC GCA GCT TTC TG-3' (down) (SEQ ID NO:)
15	pPICZ α A/mendo	5' GGG AAT TCC ATA CTC ATC AGG ACT TT-3' (up) (SEQ ID NO:)

	5' AAG CGG CCG CCT ATT TGG AGA AAG AGG T-3' (down) (SEQ ID NO:)
pPICZ α A/His.men do	5' AAG AAT TCC ATC ATC ATC ATC ATC ACA GCA GC-3' (up) (SEQ ID NO:)
	5' AAG CGG CCG CCT ATT TGG AGA AAG AGG T-3' (down) (SEQ ID NO:)
pPICZ α A/Hendo	5' TTT GAA TTC GCC CAC AGC CAC CGC GAC TTC CAG CCG GTG CTC CA-3' (up) (SEQ ID NO:)
	5' AAA AGC GGC CGC CTA CTT GGA GGC AGT CAT GAA GCT GTT CTC AA-3' (down) (SEQ ID NO:)
pPICZ α A/Restin	5' TTT TTT GAA TTC ATT TCA AGT GCC AAT TAT GAG AAG CCT GCT CTG CAT-3' (up) (SEQ ID NO:)
	5' AAG AAT GCG GCC GCT TAC TTC CTA GCG TCT GTC ATG AAA CTG TTT TCG AT-3' (down) (SEQ ID NO:)
5 pPICZ α A/HIS.Res tin	5' AAT TCC ATC ACC ATC ACC ATC ACG-3' (up) (SEQ ID NO:)
	5' AAT TCG TGA TGG TGA TGG TGA TGG-3' (down) (SEQ ID NO:)
pET28a/M2	5' TTT CAT ATG ATA TAC TCC TTT GAT GGT CGA GAC ATA ATG ACA-3' (up) (SEQ ID NO:)
	5' AAT GCG GCC GCT TAC TTC CTA GCG TCT GTC ATG AAA CTG TTT TCG AT-3' (down) (SEQ ID NO:)
pPICZ α A/M2	5' AAG AAT TCC ATC ATC ATC ATC ATC ACA GCA GC-3' (up) (SEQ ID NO:)
	5' AAT GCG GCC GCT TAC TTC CTA GCG TCT GTC ATG AAA CTG TTT TCG AT-3' (down) (SEQ ID NO:)

The resulting amplification product can then be cloned into a suitable vector.
 10 The term "primer" denotes a specific oligonucleotide-sequence-complementary to a

target nucleotide sequence and used to hybridize to the target nucleotide sequence and serve as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase. "EM 1," as used herein, refers to a deletion mutant of endostatin, wherein the last nine amino acid
5 residues have been deleted (*i.e.*, NSFMTSFSK), and the term is intended to include fragments, mutants, homologs, analogs, and allelic variants of the amino acid sequence of SEQ ID NO:). Although EM 1 was originally cloned from mouse nucleic acid, it performs better than intact type endostatin (*i.e.*, endostatin that has not been mutated) in standard assays. The term EM 1 is therefore intended to
10 include any mammalian sequence substantially similar to EM 1 as described herein, as well as mammalian EM 1 fragments, mutants, homologs, analogs and allelic variants of the mammalian EM 1 amino acid sequence. Also, specifically encompassed by the present invention are human endostatin mutants, and more specifically, the human deletion mutant equivalent of EM 1.

15 It is to be understood that the present invention is contemplated to include any derivatives of EM 1 that have endothelial inhibitory activity (*e.g.*, the capability of a composition to inhibit angiogenesis in general and, for example, to inhibit the growth or migration of bovine capillary endothelial cells in culture in the presence of fibroblast growth factor, angiogenesis-associated factors, or other known growth
20 factors). The present invention includes the entire EM 1 protein, derivatives of the EM 1 protein and biologically-active fragments of the EM 1 protein. These include proteins with EM 1 activity that have amino acid substitutions or have sugars or other molecules attached to amino acid functional groups. The present invention also includes genes that code for EM 1 and the EM 1 receptor, and to proteins that
25 are expressed by those genes.

The invention also encompasses a composition comprising an isolated polynucleotide encoding EM 1, as well as vectors and host cells containing such a polynucleotide, and processes for producing EM 1 and its fragments, mutants,
homologs, analogs and allelic variants. The term "vector" as used herein means a
30 carrier into which pieces of nucleic acid may be inserted or cloned, which carrier

functions to transfer the pieces of nucleic acid into a host cell. Such a vector may also bring about the replication and/or expression of the transferred nucleic acid pieces. Examples of vectors include nucleic acid molecules derived, *e.g.*, from a plasmid, bacteriophage, or mammalian, plant or insect virus, or non-viral vectors
5 such as ligand-nucleic acid conjugates, liposomes, or lipid-nucleic acid complexes. It may be desirable that the transferred nucleic molecule is operatively linked to an expression control sequence to form an expression vector capable of expressing the transferred nucleic acid. Such transfer of nucleic acids is generally called "transformation," and refers to the insertion of an exogenous polynucleotide into a
10 host cell, irrespective of the method used for the insertion. For example, direct uptake, transduction or f-mating are included. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome. "Operably linked" refers to a situation wherein the components described are in a relationship permitting them to function
15 in their intended manner, *e.g.*, a control sequence "operably linked" to a coding sequence is ligated in such a manner that expression of the coding sequence is achieved under conditions compatible with the control sequence. A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and translated into a polypeptide when placed under the control of (*e.g.*, operably linked
20 to) appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. Such boundaries can be naturally-occurring, or can be introduced into or added the polynucleotide sequence by methods known in the art. A coding sequence can include, but is not limited to, mRNA, cDNA, and
25 recombinant polynucleotide sequences.

The vector into which the cloned polynucleotide is cloned may be chosen because it functions in a prokaryotic organism, or alternatively, it is chosen because it functions in a eukaryotic organism. Two examples of vectors which allow for both the cloning of a polynucleotide encoding the EM 1 protein, and the expression of
30 that protein from the polynucleotide are the pET28(a) vector (Novagen, Madison,

Wisconsin, USA) and a modified pPICZ α A vector (Invitrogen, San Diego, California, USA), which allow expression of the protein in bacteria and yeast, respectively.

Once a polynucleotide has been cloned into a suitable vector, it can be transformed into an appropriate host cell. By "host cell" is meant a cell which has been or can be used as the recipient of transferred nucleic acid by means of a vector. Host cells can be prokaryotic or eukaryotic, mammalian, plant, or insect, and can exist as single cells, or as a collection, *e.g.*, as a culture, or in a tissue culture, or in a tissue or an organism. Host cells can also be derived from normal or diseased tissue from a multicellular organism, *e.g.*, a mammal. Host cell, as used herein, is intended to include not only the original cell which was transformed with a nucleic acid, but also descendants of such a cell, which still contain the nucleic acid.

In one embodiment, the isolated polynucleotide encoding the anti-angiogenic protein additionally comprises a polynucleotide linker encoding a peptide. Such linkers are known to those of skill in the art and, for example the linker can comprise at least one additional codon encoding at least one additional amino acid. Typically the linker comprises one to about twenty or thirty amino acids. The polynucleotide linker is translated, as is the polynucleotide encoding the anti-angiogenic protein, resulting in the expression of an anti-angiogenic protein with at least one additional amino acid residue at the amino or carboxyl terminus of the anti-angiogenic protein. Some linkers attached to anti-angiogenic proteins are illustrated in Figure 26. Importantly, the additional amino acid, or amino acids, do not compromise the activity of the anti-angiogenic protein.

After inserting the selected polynucleotide into the vector, the vector is transformed into an appropriate prokaryotic strain and the strain is cultured (*e.g.*, maintained) under suitable culture conditions for the production of the biologically active anti-angiogenic protein, thereby producing a biologically active anti-angiogenic protein, or mutant, derivative, fragment or fusion protein thereof. In one embodiment, the invention comprises cloning of a polynucleotide encoding an anti-angiogenic protein into the vectors pET17b or pET28a, which are then transformed into bacteria. The bacterial host strain then expresses the anti-

angiogenic protein. Typically the anti-angiogenic proteins are produced in quantities of about 10-20 milligrams. or more, per liter of culture fluid.

In another embodiment of the present invention, the eukaryotic vector comprises a yeast vector. As described herein, one method uses a pPICZ α plasmid wherein the plasmid contains a multiple cloning site. The multiple cloning site inserted into the multiple cloning site a His.Tag motif. Additionally the vector can be modified to add a *NdeI* site, or other suitable restriction sites. Such sites are well known to those of skill in the art. Anti-angiogenic proteins produced by this embodiment comprise a histidine tag motif (His.tag) comprising one, or more histidines, typically about 5-20 histidines. Surprisingly, this His.tag does not compromise anti-angiogenic activity.

In this embodiment, a preferred yeast expression system is *Pichia pastores*. Again, the biologically active protein is typically produced at concentrations of about 10-20 milligrams per liter of culture medium (fluid).

One method of producing EM 1, for example, is to amplify the polynucleotide of SEQ ID NO:1, clone it into an expression vector, *e.g.*, pET28(a), pPICZ α A, or some other expression vector, transform the vector containing the polynucleotide of SEQ ID NO:1 into a host cell capable of expressing the polypeptide encoded by the polynucleotide, culturing the transformed host cell under culture conditions suitable for expressing the protein, and then extracting and purifying the protein from the culture. Exemplary methods of producing anti-angiogenic proteins in general, and EM 1 in particular, are provided in the Examples below, and also in U.S.S.N. XX/XXX,XXX, "Methods of Producing Anti-Angiogenic Proteins," by Vikas P. Sukhatme, filed December 8, 1998, the entire teachings of all of which are herein incorporated by reference. The EM 1 protein may also be expressed as a product of transgenic animals, *e.g.*, as a component of the milk of transgenic cows, goats, sheep or pigs, or as a product of a transgenic plant, *e.g.*, combined or linked with starch molecules in maize.

EM 1 may also be produced by conventional, known methods of chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed EM 1

protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with *e.g.*, recombinantly-produced EM 1, may possess biological properties in common therewith, including biological activity. Thus, the synthetically-constructed EM 1 protein sequences may be employed as

5 biologically active or immunological substitutes for *e.g.*, recombinantly-produced, purified EM 1 protein in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The EM 1 protein is useful in inhibiting angiogenesis, as determined in standard assays, and provided in the Examples below. EM 1 does not inhibit the

10 growth of other cells types, *e.g.*, IMR-90 cells, or IC-21 cells.

As used herein, the term "angiogenesis" means the generation of new blood vessels into a tissue or organ, and involves endothelial cell proliferation. Under normal physiological conditions, humans or animals undergo angiogenesis only in very specific restricted situations. For example, angiogenesis is normally observed

15 in wound healing, fetal and embryonal development, and formation of the corpus luteum, endometrium and placenta. The term "endothelium" means a thin layer of flat epithelial cells that lines serous cavities, lymph vessels, and blood vessels. "Anti-angiogenic activity" therefore refers to the capability of a composition to inhibit the growth of blood vessels. The growth of blood vessels is a complex series

20 of events, and includes localized breakdown of the basement membrane lying under the individual endothelial cells, proliferation of those cells, migration of the cells to the location of the future blood vessel, reorganization of the cells to form a new vessel membrane, cessation of endothelial cell proliferation, and, incorporation of pericytes and other cells that support the new blood vessel wall. "Anti-angiogenic

25 activity" as used herein therefore includes interruption of any or all of these stages, with the end result that formation of new blood vessels is inhibited.

Anti-angiogenic activity may include endothelial inhibiting activity, which refers to the capability of a composition to inhibit angiogenesis in general and, for example, to inhibit the growth or migration of bovine capillary endothelial cells in

30 culture in the presence of fibroblast growth factor, angiogenesis-associated factors,

or other known growth factors. A "growth factor" is a composition that stimulates the growth, reproduction, or synthetic activity of cells. An "angiogenesis-associated factor" is a factor which either inhibits or promotes angiogenesis. An example of an angiogenesis-associated factor is an angiogenic growth factor, such as basic fibroblastic growth factor (bFGF), which is an angiogenesis promoter. Another example of an angiogenesis-associated factor is an angiogenesis inhibiting factor such as *e.g.*, angiostatin (see, *e.g.*, U.S. Pat. No. 5,801,012, U.S. Pat. No. 5,837,682, U.S. Pat. No. 5,733,876, U.S. Pat. No. 5,776,704, U.S. Pat. No. 5,639,725, U.S. Pat. No. 5,792,845, WO 96/35774, WO 95/29242, WO 96/41194, WO 97/23500) or endostatin (see, *e.g.*, WO 97/15666).

By "substantially the same biological activity" or "substantially the same or superior biological activity" is meant that a composition has anti-angiogenic activity, and behaves similarly as does EM 1, as determined in standard assays. "Standard assays" include, but are not limited to, those protocols used in the molecular biological arts to assess anti-angiogenic activity, cell cycle arrest, and apoptosis. Such assays include, but are not limited to, assays of endothelial cell proliferation, endothelial cell migration, cell cycle analysis, and endothelial cell tube formation, detection of apoptosis, *e.g.*, by apoptotic cell morphology or Annexin V-FITC assay, chorioallantoic membrane (CAM) assay, and inhibition of renal cancer tumor growth in nude mice. Such assays are provided in the Examples below, and in U.S.S.N. 60/067,888, filed December 8, 1997, U.S.S.N. 60/082,663, filed April 22, 1998, U.S.S.N. 60/108,536, filed November 16, 1998, and in U.S.S.N. XX/XXX,XXX, "Restin and Methods of Use Thereof," by Vikas P. Sukhatme, filed December 8, 1998, and U.S.S.N. XX/XXX,XXX, "Methods of Producing Anti-Angiogenic Proteins," by Vikas P. Sukhatme, filed December 8, 1998, the entire teachings of all of which are herein incorporated by reference. Such methods are also included in Dhanabal *et al.* (1998) ("Endostatin Induces Endothelial Cell Apoptosis," *J. Biol. Chem.*, submitted), and in Dhanabal *et al.* (1999) ("Cloning, Expression and *in vitro* Activity of Human Endostatin," *Cancer Res.*, in press). Evaluating the ED₅₀ of a

mutant in one of the assays described herein is a useful method of comparing activities.

As used herein, "ED₅₀" is an abbreviation for the amount of a composition which reduces a biological effect by one-half, relative to the biological effect seen in the absence of the composition.

The invention also describes fragments, mutants, homologs and analogs of EM 1. A "fragment" of EM 1 any amino acid sequence shorter than the EM 1 molecule, comprising at least 25 consecutive amino acids of the EM 1 polypeptide. Such mutants may or may not also comprise additional amino acids derived from the process of cloning, *e.g.*, amino acid residues or amino acid sequences corresponding to full or partial linker sequences. To be encompassed by the present invention, such mutants, with or without such additional amino acid residues, must have substantially the same biological activity as the natural or full-length version of the reference polypeptide.

By "mutant" of EM 1 is meant a polypeptide that includes any change in the amino acid sequence relative to the amino acid sequence of the equivalent reference EM 1 polypeptide. Such changes can arise either spontaneously or by manipulations by man, by chemical energy (*e.g.*, X-ray), or by other forms of chemical mutagenesis, or by genetic engineering, or as a result of mating or other forms of exchange of genetic information. Mutations include, *e.g.*, base changes, deletions, insertions, inversions, translocations, or duplications. Mutant forms of EM 1 may display either increased or decreased anti-angiogenic activity relative to the equivalent reference EM 1 polynucleotide, and such mutants may or may not also comprise additional amino acids derived from the process of cloning, *e.g.*, amino acid residues or amino acid sequences corresponding to full or partial linker sequences.

By "analog" of EM 1 is meant a non-natural molecule substantially similar to either the entire EM 1 molecule or a fragment or allelic variant thereof, and having substantially the same or superior biological activity. Such analogs are intended to include derivatives (*e.g.*, chemical derivatives, as defined above) of the biologically

active EM 1, as well as its fragments, mutants, homologs, and allelic variants, which derivatives exhibit a qualitatively similar agonist or antagonist effect to that of the unmodified EM 1 polypeptide, fragment, mutant, homolog, or allelic variant.

By "allele" of EM 1 is meant a polypeptide sequence containing a naturally-
5 occurring sequence variation relative to the polypeptide sequence of the reference EM 1 polypeptide. By "allele" of a polynucleotide encoding the EM 1 polypeptide is meant a polynucleotide containing a sequence variation relative to the reference polynucleotide sequence encoding the reference EM 1 polypeptide, where the allele of the polynucleotide encoding the EM 1 polypeptide encodes an allelic form of the
10 EM 1 polypeptide.

It is possible that a given polypeptide may be either a fragment, a mutant, an analog, or allelic variant of EM 1, or it may be two or more of those things, *e.g.*, a polypeptide may be both an analog and a mutant of the EM 1 polypeptide. For example, a shortened version of the EM 1 molecule (*e.g.*, a fragment of EM 1) may
15 be created in the laboratory. If that fragment is then mutated through means known in the art, a molecule is created that is both a fragment and a mutant of EM 1. In another example, a mutant of EM 1 may be created, which is later discovered to exist as an allelic of EM 1 in some mammalian individuals. Such a mutant EM 1 molecule would therefore be both a mutant and an allelic variant of EM 1. Such
20 combinations of fragments, mutants, allelic variants, and analogs are intended to be encompassed in the present invention.

Encompassed by the present invention are proteins that have substantially the same amino acid sequence as EM 1, or polynucleotides that have substantially the same nucleic acid sequence as the polynucleotide encoding EM 1. "Substantially the
25 same sequence" means a nucleic acid or polypeptide that exhibits at least about 70 % sequence identity with a reference sequence, *e.g.*, another nucleic acid or polypeptide, typically at least about 80% sequence identity with the reference sequence, preferably at least about 90% sequence identity, more preferably at least about 95% identity, and most preferably at least about 97% sequence identity with
30 the reference sequence. The length of comparison for sequences will generally be at

least 75 nucleotide bases or 25 amino acids, more preferably at least 150 nucleotide bases or 50 amino acids, and most preferably 243-264 nucleotide bases or 81-88 amino acids. "Polypeptide" as used herein indicates a molecular chain of amino acids and does not refer to a specific length of the product. Thus, peptides, oligopeptides and proteins are included within the definition of polypeptide. This term is also intended to include polypeptide that have been subjected to post-expression modifications such as, for example, glycosylations, acetylations, phosphorylations and the like. EM 1, in general, has less than 70% amino acid sequence identity with endostatin.

"Sequence identity," as used herein, refers to the subunit sequence similarity between two polymeric molecules, *e.g.*, two polynucleotides or two polypeptides. When a subunit position in both of the two molecules is occupied by the same monomeric subunit, *e.g.*, if a position in each of two peptides is occupied by serine, then they are identical at that position. The identity between two sequences is a direct function of the number of matching or identical positions, *e.g.*, if half (*e.g.*, 5 positions in a polymer 10 subunits in length), of the positions in two peptide or compound sequences are identical, then the two sequences are 50% identical; if 90% of the positions, *e.g.*, 9 of 10 are matched, the two sequences share 90% sequence identity. By way of example, the amino acid sequences VRGLQP and HAFLQP have 3 of 6 positions in common, and therefore share 50% sequence identity, while the sequences VRGLQP and AFLQP have 3 of 5 positions in common, and therefore share 60% sequence identity. The identity between two sequences is a direct function of the number of matching or identical positions. Thus, if a portion of the reference sequence is deleted in a particular peptide, that deleted section is not counted for purposes of calculating sequence identity, *e.g.*, VRGLQP and VRGLP have 5 out of 6 position in common, and therefore share 83.3% sequence identity.

Identity is often measured using sequence analysis software *e.g.*, BLASTN or BLASTP (available at <http://www.ncbi.nlm.nih.gov/BLAST/>). The default parameters for comparing two sequences (*e.g.*, "Blast"-ing two sequences against each other, <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>) by BLASTN (for nucleotide

sequences) are reward for match = 1, penalty for mismatch = -2, open gap = 5, extension gap = 2. When using BLASTP for protein sequences, the default parameters are reward for match = 0, penalty for mismatch = 0, open gap = 11, and extension gap = 1.

5 When two sequences share "sequence homology," it is meant that the two sequences differ from each other only by conservative substitutions. For polypeptide sequences, such conservative substitutions consist of substitution of one amino acid at a given position in the sequence for another amino acid of the same class (*e.g.*, amino acids that share characteristics of hydrophobicity, charge, pK or
10 other conformational or chemical properties, *e.g.*, valine for leucine, arginine for lysine), or by one or more non-conservative amino acid substitutions, deletions, or insertions, located at positions of the sequence that do not alter the conformation or folding of the polypeptide to the extent that the biological activity of the polypeptide is destroyed. Examples of "conservative substitutions" include substitution of one
15 non-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another; the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine; the substitution of one basic residue such as lysine, arginine or histidine for another; or the substitution of one acidic residue, such as aspartic acid or
20 glutamic acid for another; or the use of a chemically derivatized residue in place of a non-derivatized residue; provided that the polypeptide displays the requisite biological activity. Two sequences which share sequence homology may be called "sequence homologs."

Homology, for polypeptides, is typically measured using sequence analysis
25 software (*e.g.*, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). Protein analysis software matches similar sequences by assigning degrees of homology to various substitutions, deletions, and other modifications. Conservative substitutions typically include substitutions within the
30 following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid,

glutamic acid, asparagine, glutamine; serine threonine; lysine, arginine; and phenylalanine, tyrosine.

Also encompassed by the present invention are chemical derivatives of EM 1. "Chemical derivative" refers to a subject polypeptide having one or more residues chemically derivatized by reaction of a functional side group. Such
5 derivatized residues include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and
10 ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-imbenzylhistidine. Also included as chemical derivatives are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids. For examples:
15 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine.

Polynucleotides encoding EM 1 can be cloned out of isolated DNA or a cDNA library. Nucleic acids polypeptides, referred to herein as "isolated" are
20 nucleic acids or polypeptides substantially free (*i.e.*, separated away from) the material of the biological source from which they were obtained (*e.g.*, as exists in a mixture of nucleic acids or in cells), which may have undergone further processing. "Isolated" nucleic acids or polypeptides include nucleic acids or polypeptides obtained by methods described herein, similar methods, or other suitable methods,
25 including essentially pure nucleic acids or polypeptides, nucleic acids or polypeptides produced by chemical synthesis, by combinations of chemical or biological methods, and recombinantly produced nucleic acids or polypeptides which are isolated. An isolated polypeptide therefore means one which is relatively free of other proteins, carbohydrates, lipids, and other cellular components with which it is
30 normally associated. An isolated nucleic acid is not immediately contiguous with

(i.e., covalently linked to) both of the nucleic acids with which it is immediately contiguous in the naturally-occurring genome of the organism from which the nucleic acid is derived. The term, therefore, includes, for example, a nucleic acid which is incorporated into a vector (e.g., an autonomously replicating virus or
5 plasmid), or a nucleic acid which exists as a separate molecule independent of other nucleic acids such as a nucleic acid fragment produced by chemical means or restriction endonuclease treatment.

The polynucleotides and proteins of the present invention can also be used to design probes to isolate other anti-angiogenic proteins. Exceptional methods are
10 provided in U.S. Pat. No. 5,837,490, by Jacobs *et al.*, the entire teachings of which are herein incorporated by reference in their entirety. The design of the oligonucleotide probe should preferably follow these parameters: (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any, and (b) It should be designed to have a T_m of approx. 80° C (assuming 2° C for
15 each A or T and 4 degrees for each G or C).

The oligonucleotide should preferably be labeled with γ - ^{32}P ATP (specific activity 6000 Ci/mmmole) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography
20 or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a scintillation counter. Preferably, specific activity of the resulting probe should be approximately 4×10^6 dpm/pmmole. The bacterial culture containing the pool of full-length clones should preferably be thawed and 100 μl of the stock used to inoculate a sterile culture flask containing 25
25 ml of sterile L-broth containing ampicillin at 100 $\mu\text{g}/\text{ml}$. The culture should preferably be grown to saturation at 37°C, and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing
30 L-broth containing ampicillin at 100 $\mu\text{g}/\text{ml}$ and agar at 1.5% in a 150 mm petri dish

when grown overnight at 37°C. Other known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them. Highly stringent
5 condition are those that are at least as stringent as, for example, 1x SSC at 65°C, or 1x SSC and 50% formamide at 42°C. Moderate stringency conditions are those that are at least as stringent as 4x SSC at 65°C, or 4x SSC and 50% formamide at 42°C. Reduced stringency conditions are those that are at least as stringent as 4x SSC at 50°C, or 6x SSC and 50% formamide at 40°C.

10 The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6.times. SSC (20x stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100 µg/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is
15 10⁶ dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2x SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2x SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with
20 0.1x SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed. The positive colonies are then picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by
restriction analysis, hybridization analysis, or DNA sequencing.

25 The present invention also includes fusion proteins and chimeric proteins comprising EM 1, its fragments, mutants, homologs, analogs, and allelic variants. A fusion or chimeric protein can consist of a multimer of a single protein, *e.g.*, repeats of EM 1 or repeats of apomigren, or the fusion and chimeric proteins can be made up of several proteins, *e.g.*, EM 1 and apomigren. The fusion proteins can comprise a
30 combination of two or more known anti-angiogenic proteins (*e.g.*, angiostatin,

endostatin, restin, or apomigren, or biologically active fragments thereof), or an anti-angiogenic protein in combination with a targeting agent (*e.g.*, endostatin with epidermal growth factor (EGF) or RGD peptides), or an anti-angiogenic protein in combination with an immunoglobulin molecule (*e.g.*, endostatin and IgG,
5 specifically with the Fc portion removed). As used herein, "restin" is a protein comprising about 170 to about 200 amino acid residues, and has at least 70% sequence identity with the C-terminus of the NC10 domain of the $\alpha 1$ chain of human Type XV collagen. As used herein, "apomigren" is a fragment of restin, and comprises the last 80 to 90 contiguous amino acids corresponding to the C-terminus
10 of the NC10 domain of the $\alpha 1$ chain of human Type XV collagen. The fusion and chimeric proteins can also include EM 1, its fragments, mutants, homologs, analogs, and allelic variants, and other anti-angiogenic proteins, *e.g.*, endostatin or angiostatin. The term "fusion protein" as used herein can also encompass additional components for *e.g.*, delivering a chemotherapeutic agent, wherein a polynucleotide
15 encoding the chemotherapeutic agent is linked to the polynucleotide encoding the anti-angiogenic protein. Fusion proteins can also encompass multimers of the anti-angiogenic protein, *e.g.*, a dimer or trimer of endostatin. Such fusion proteins can be linked together via post-translational modification (*e.g.*, chemically linked), or the entire fusion protein may be made recombinantly.

20 Also included in the inventions are compositions containing, as a biological ingredient, EM 1, as well as its fragments, mutants, homologs, analogs, and allelic variants to inhibit or enhance angiogenesis in mammalian tissues, and use of such compositions in the diagnosis, prognosis, and treatment of diseases and conditions characterized by, or associated with, angiogenic activity or lack thereof. Such
25 methods can involve administration by oral, topical, injection, implantation, sustained release, or other delivery methods.

The invention includes use of EM 1, and its fragments, mutants, homologs, analogs, allelic variants, and fusion and chimeric proteins as biologically-active agents in compositions for the purpose of treating diseases or conditions that are
30 associated with angiogenic activity. Methods of treating such diseases include

contacting the affected tissue with a composition comprising EM 1, its fragments, mutants, homologs, analogs, or allelic variants.

The present invention includes the method of treating an angiogenesis-mediated disease with a therapeutically effective amount of EM 1, or a biologically
5 active fragment thereof, or combinations of EM 1 fragments that possess anti-angiogenic activity, or EM 1 agonists and antagonists. Angiogenesis-mediated diseases include, but are not limited to, cancers, solid tumors, blood-born tumors (*e.g.*, leukemias), tumor metastasis, benign tumors (*e.g.*, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas), rheumatoid
10 arthritis, psoriasis, ocular angiogenic diseases (*e.g.*, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis), Osler-Webber Syndrome, myocardial angiogenesis, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma, and wound granulation. EM 1 is useful in the treatment of diseases of
15 excessive or abnormal stimulation of endothelial cells. These diseases include, but are not limited to, intestinal adhesions, Crohn's disease, atherosclerosis, scleroderma, and hypertrophic scars (*i.e.*, keloids). EM 1 can be used as a birth control agent by preventing vascularization required for embryo implantation. EM 1 is useful in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat
20 scratch disease (*Rochele minalia quintosa*) and ulcers (*Helicobacter pylori*). EM 1 can also be used to prevent dialysis graft vascular access stenosis, and obesity, *e.g.*, by inhibiting capillary formation in adipose tissue, thereby preventing its expansion. EM 1 can also be used to treat localized (*e.g.*, nonmetastisized) diseases. "Cancer" means neoplastic growth, hyperplastic or proliferative growth or a pathological state
25 of abnormal cellular development and includes solid tumors, non-solid tumors, and any abnormal cellular proliferation, such as that seen in leukemia. As used herein, "cancer" also means angiogenesis-dependent cancers and tumors, *i.e.*, tumors that require for their growth (expansion in volume and/or mass) an increase in the number and density of the blood vessels supplying them with blood. "Regression"
30 refers to the reduction of tumor mass and size. As used herein, the term

"therapeutically effective amount" means the total amount of each active component of the composition or method that is sufficient to show a meaningful patient benefit, *i.e.*, treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such

5 conditions. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

Alternatively, where an increase in angiogenesis is desired, *e.g.*, in wound healing, or in post-infarct heart tissue, antibodies or antisera to the EM 1 protein can

10 be used to block localized, native anti-angiogenic proteins and processes, and thereby increase formation of new blood vessels so as to inhibit atrophy of tissue.

EM 1 may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation, chemotherapy, or immunotherapy combined with EM 1 and

15 then EM 1 may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor. EM 1, EM 1 fragments, EM 1 antisera, EM 1 receptor agonists, EM 1 receptor antagonists, or combinations thereof, can also be combined with other anti-angiogenic compounds, or proteins, fragments, antisera, receptor agonists, receptor

20 antagonists of other anti-angiogenic proteins (*e.g.*, angiostatin, endostatin, restin, apomigren). Additionally, EM 1, EM 1 fragments, EM 1 antisera, EM 1 receptor agonists, EM 1 receptor antagonists, or combinations thereof, are combined with pharmaceutically acceptable excipients, and optionally sustained-release matrix, such as biodegradable polymers, to form therapeutic compositions. The

25 compositions of the present invention may also contain other anti-angiogenic proteins or chemical compounds, such as endostatin, angiostatin, restin and apomigren (both of which are described in in U.S.S.N. XX/XXX,XXX, "Restin and Methods of use Thereof", by Vikas P. Sukhatme, filed December 8, 1998, the entire teachings of which are herein incorporated by reference), and mutants, fragments,

30 and analogs thereof. The compositions may further contain other agents which

either enhance the activity of the protein or compliment its activity or use in treatment, such as chemotherapeutic or radioactive agents. Such additional factors and/or agents may be included in the composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Additionally,

5 administration of the composition of the present invention may be administered concurrently with other therapies, *e.g.*, administered in conjunction with a chemotherapy or radiation therapy regimen.

The invention includes methods for inhibiting angiogenesis in mammalian tissues by contacting the tissue with a composition comprising the proteins of the invention. By "contacting" is meant not only topical application, but also those

10 modes of delivery that introduce the composition into the tissues, or into the cells of the tissues.

Use of timed release or sustained release delivery systems are also included in the invention. Such systems are highly desirable in situations where surgery is

15 difficult or impossible, *e.g.*, patients debilitated by age or the disease course itself, or where the risk-benefit analysis dictates control over cure.

A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid/base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and

20 body fluids. The sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (co-polymers of lactic acid and glycolic acid) polyanhydrides, poly(ortho)esters, polyproteins, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides,

25 nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

The angiogenesis-modulating therapeutic composition of the present invention may be a solid, liquid or aerosol and may be administered by any known route of administration. Examples of solid compositions include pills, creams, and implantable dosage units. The pills may be administered orally, the creams may be administered topically. The implantable dosage unit may be administered locally, for example at a tumor site, or which may be implanted for systemic release of the angiogenesis-modulating composition, for example subcutaneously. Examples of liquid composition include formulations adapted for injection subcutaneously, intravenously, intraarterially, and formulations for topical and intraocular administration. Examples of aerosol formulation include inhaler formulation for administration to the lungs.

The EM 1 proteins and protein fragments with the anti-angiogenic activity described above can be provided as isolated and substantially purified proteins and protein fragments in pharmaceutically acceptable formulations using formulation methods known to those of ordinary skill in the art. These formulations can be administered by standard routes. In general, the combinations may be administered by the topical, transdermal, intraperitoneal, intracranial, intracerebroventricular, intracerebral, intravaginal, intrauterine, oral, rectal or parenteral (*e.g.*, intravenous, intraspinal, subcutaneous or intramuscular) route. In addition, the EM 1 may be incorporated into biodegradable polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the EM 1 is slowly released systemically. Osmotic minipumps may also be used to provide controlled delivery of high concentrations of EM 1 through cannulae to the site of interest, such as directly into a metastatic growth or into the vascular supply to that tumor. The biodegradable polymers and their use are described, for example, in detail in Brem *et al.* (1991) (*J. Neurosurg.* 74:441-446), which is hereby incorporated by reference in its entirety.

The compositions containing a polypeptide of this invention can be administered intravenously, as by injection of a unit dose, for example. The term

"unit dose" when used in reference to a therapeutic composition of the present invention refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent; *i.e.*,
5 carrier or vehicle.

Modes of administration of the compositions of the present inventions include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion. Pharmaceutical compositions for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous
10 solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (*e.g.*, glycerol, propylene glycol, polyethylene glycol and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils
15 (*e.g.*, olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention
20 of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents such as paraben, chlorobutanol, phenol sorbic acid and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents, such as aluminum
25 monostearate and gelatin, which delay absorption. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters) and poly(anhydrides). Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Depot injectable formulations are also
30 prepared by entrapping the drug in liposomes or microemulsions which are

compatible with body tissues. The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable media just prior to use.

5 The therapeutic compositions of the present invention can include pharmaceutically acceptable salts of the components therein, *e.g.*, which may be derived from inorganic or organic acids. By "pharmaceutically acceptable salt" is meant those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue
10 toxicity, irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well-known in the art. For example, S. M. Berge, et al. describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences* (1977) 66:1 *et seq.*, which is incorporated herein by reference. Pharmaceutically acceptable salts include the acid addition salts
15 (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium or ferric hydroxides, and such organic
20 bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like. The salts may be prepared *in situ* during the final isolation and purification of the compounds of the invention or separately by reacting a free base function with a suitable organic acid. Representative acid addition salts include, but are not limited to acetate, adipate, alginate, citrate, aspartate, benzoate,
25 benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxymethanesulfonate (isethionate), lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate,
30 tartate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and

undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; arylalkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained. Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, hydrobromic acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

As used herein, the terms "pharmaceutically acceptable," "physiologically tolerable" and grammatical variations thereof as they refer to compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are capable of administration to or upon a mammal with a minimum of undesirable physiological effects such as nausea, dizziness, gastric upset and the like. The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art and need not be limited based on formulation. Typically such compositions are prepared as injectables either as liquid solutions or suspensions, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified.

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients include, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The EM 1 polypeptides of the present invention can also be included in a composition comprising a prodrug. As used herein, the term "prodrug" refers to

compounds which are rapidly transformed *in vivo* to yield the parent compound, for example, by enzymatic hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, *Prodrugs as Novel Delivery Systems*, Vol. 14 of the ACS Symposium Series and in Edward B. Roche, ed., *Bioreversible Carriers in Drug*
5 *Design*, American Pharmaceutical Association and Permagon Press, 1987, both of which are incorporated herein by reference. As used herein, the term "pharmaceutically acceptable prodrug" refers to (1) those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and animals without undue
10 toxicity, irritation, allergic response and the like, commensurate with a suitable benefit-to-risk ratio and effective for their intended use and (2) zwitterionic forms, where possible, of the parent compound.

The dosage of the EM 1 of the present invention will depend on the disease state or condition being treated and other clinical factors such as weight and
15 condition of the human or animal and the route of administration of the compound. For treating humans or animals, about 10 mg/kg of body weight to about 20 mg/kg of body weight of the EM 1 protein or the apomigren protein can be administered. In combination therapies, *e.g.*, the EM 1 protein of the invention in combination with radiotherapy, chemotherapy, or immunotherapy, it may be possible to reduce
20 the dosage, *e.g.*, to about 0.1 mg/kg of body weight to about 0.2 mg/kg of body weight. Depending upon the half-life of the EM 1 in the particular animal or human, the EM 1 can be administered between several times per day to once a week. It is to be understood that the present invention has application for both human and veterinary use. The methods of the present invention contemplate single as well as
25 multiple administrations, given either simultaneously or over an extended period of time. In addition, EM 1 can be administered in conjunction with other forms of therapy, *e.g.*, chemotherapy, radiotherapy, or immunotherapy.

The EM 1 formulations include those suitable for oral, rectal, ophthalmic (including intravitreal or intracameral), nasal, topical (including buccal and
30 sublingual), intrauterine, vaginal or parenteral (including subcutaneous,

intraperitoneal, intramuscular, intravenous, intradermal, intracranial, intratracheal, and epidural) administration. The EM 1 formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

When a therapeutically effective amount of protein of the present invention is administered orally, the EM 1 protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene

glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

5 When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or
10 subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of
15 the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

 The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has
20 undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at
25 that point the dosage is not increased further.

 The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the
30 present invention will be in the range of 12 to 24 hours of continuous intravenous

administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly mentioned above, the formulations of the present invention may include other agents conventional in the art having regard to the type of formulation in question. Optionally, cytotoxic agents may be incorporated or otherwise combined with EM 1 proteins, or biologically functional protein fragments thereof, to provide dual therapy to the patient.

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins of the present invention.

Cytotoxic agents such as ricin, are linked to EM 1, and high affinity EM 1 protein fragments, thereby providing a tool for destruction of cells that bind EM 1. These cells may be found in many locations, including but not limited to, micrometastases and primary tumors. Proteins linked to cytotoxic agents are infused in a manner designed to maximize delivery to the desired location. For example, ricin-linked high affinity EM 1 fragments are delivered through a cannula into vessels supplying the target site or directly into the target. Such agents are also delivered in a controlled manner through osmotic pumps coupled to infusion cannulae. A combination of EM 1 antagonists may be co-applied with stimulators of angiogenesis to increase vascularization of tissue. This therapeutic regimen provides an effective means of destroying metastatic cancer.

Additional treatment methods include administration of EM 1, EM 1 fragments, EM 1 analogs, EM 1 antisera, or EM 1 receptor agonists and antagonists linked to cytotoxic agents. It is to be understood that the EM 1 can be human or animal in origin. EM 1 can also be produced synthetically by chemical reaction or

by recombinant techniques in conjunction with expression systems. EM 1 can also be produced by enzymatically cleaving isolated plasminogen or plasmin to generate proteins having anti-angiogenic activity. EM 1 may also be produced by compounds that mimic the action of endogenous enzymes that cleave plasminogen to EM 1.

5 EM 1 production may also be modulated by compounds that affect the activity of plasminogen cleaving enzymes.

The present invention also encompasses gene therapy whereby a polynucleotide encoding EM 1, or a mutant, fragment, or fusion protein thereof, is introduced and regulated in a patient. Various methods of transferring or delivering
10 DNA to cells for expression of the gene product protein, otherwise referred to as gene therapy, are disclosed in *Gene Transfer into Mammalian Somatic Cells in vivo*, N. Yang (1992) *Crit. Rev. Biotechn.* 12(4):335-356, which is hereby incorporated by reference. Gene therapy encompasses incorporation of DNA sequences into somatic cells or germ line cells for use in either ex vivo or in vivo therapy. Gene therapy
15 functions to replace genes, augment normal or abnormal gene function, and to combat infectious diseases and other pathologies.

Strategies for treating these medical problems with gene therapy include therapeutic strategies such as identifying the defective gene and then adding a functional gene to either replace the function of the defective gene or to augment a
20 slightly functional gene; or prophylactic strategies, such as adding a gene for the product protein that will treat the condition or that will make the tissue or organ more susceptible to a treatment regimen. As an example of a prophylactic strategy, a gene such as EM 1 may be placed in a patient and thus prevent occurrence of angiogenesis; or a gene that makes tumor cells more susceptible to radiation could
25 be inserted and then radiation of the tumor would cause increased killing of the tumor cells.

Many protocols for transfer of EM 1 DNA or EM 1 regulatory sequences are envisioned in this invention. Transfection of promoter sequences, other than one normally found specifically associated with EM 1, or other sequences which would
30 increase production of EM 1 protein are also envisioned as methods of gene therapy.

An example of this technology is found in Transkaryotic Therapies, Inc., of Cambridge, Mass., using homologous recombination to insert a "genetic switch" that turns on an erythropoietin gene in cells. See *Genetic Engineering News*, Apr. 15, 1994. Such "genetic switches" could be used to activate EM 1 (or the EM 1 receptor) in cells not normally expressing EM 1 (or the EM 1 receptor).

Gene transfer methods for gene therapy fall into three broad categories: physical (e.g., electroporation, direct gene transfer and particle bombardment), chemical (e.g., lipid-based carriers, or other non-viral vectors) and biological (e.g., virus-derived vector and receptor uptake). For example, non-viral vectors may be used which include liposomes coated with DNA. Such liposome/DNA complexes may be directly injected intravenously into the patient. It is believed that the liposome/DNA complexes are concentrated in the liver where they deliver the DNA to macrophages and Kupffer cells. These cells are long lived and thus provide long term expression of the delivered DNA. Additionally, vectors or the "naked" DNA of the gene may be directly injected into the desired organ, tissue or tumor for targeted delivery of the therapeutic DNA.

Gene therapy methodologies can also be described by delivery site. Fundamental ways to deliver genes include *ex vivo* gene transfer, *in vivo* gene transfer, and *in vitro* gene transfer. In *ex vivo* gene transfer, cells are taken from the patient and grown in cell culture. The DNA is transfected into the cells, the transfected cells are expanded in number and then reimplanted in the patient. In *in vitro* gene transfer, the transformed cells are cells growing in culture, such as tissue culture cells, and not particular cells from a particular patient. These "laboratory cells" are transfected, the transfected cells are selected and expanded for either implantation into a patient or for other uses.

In vivo gene transfer involves introducing the DNA into the cells of the patient when the cells are within the patient. Methods include using virally mediated gene transfer using a noninfectious virus to deliver the gene in the patient or injecting naked DNA into a site in the patient and the DNA is taken up by a percentage of cells in which the gene product protein is expressed. Additionally, the

other methods described herein, such as use of a "gene gun," may be used for *in vitro* insertion of EM 1 DNA or EM 1 regulatory sequences.

Chemical methods of gene therapy may involve a lipid based compound, not necessarily a liposome, to transfer the DNA across the cell membrane. Lipofectins
5 or cytofectins, lipid-based positive ions that bind to negatively charged DNA, make a complex that can cross the cell membrane and provide the DNA into the interior of the cell. Another chemical method uses receptor-based endocytosis, which involves binding a specific ligand to a cell surface receptor and enveloping and transporting it
10 across the cell membrane. The ligand binds to the DNA and the whole complex is transported into the cell. The ligand gene complex is injected into the blood stream and then target cells that have the receptor will specifically bind the ligand and transport the ligand-DNA complex into the cell.

Many gene therapy methodologies employ viral vectors to insert genes into cells. For example, altered retrovirus vectors have been used in *ex vivo* methods to
15 introduce genes into peripheral and tumor-infiltrating lymphocytes, hepatocytes, epidermal cells, myocytes, or other somatic cells. These altered cells are then introduced into the patient to provide the gene product from the inserted DNA.

Viral vectors have also been used to insert genes into cells using *in vivo* protocols. To direct the tissue-specific expression of foreign genes, *cis*-acting
20 regulatory elements or promoters that are known to be tissue-specific can be used. Alternatively, this can be achieved using *in situ* delivery of DNA or viral vectors to specific anatomical sites *in vivo*. For example, gene transfer to blood vessels *in vivo* was achieved by implanting *in vitro* transduced endothelial cells in chosen sites on arterial walls. The virus infected surrounding cells which also expressed the gene
25 product. A viral vector can be delivered directly to the *in vivo* site, by a catheter for example, thus allowing only certain areas to be infected by the virus, and providing long-term, site specific gene expression. *In vivo* gene transfer using retrovirus vectors has also been demonstrated in mammary tissue and hepatic tissue by injection of the altered virus into blood vessels leading to the organs.

Viral vectors that have been used for gene therapy protocols include but are not limited to, retroviruses, other RNA viruses such as poliovirus or Sindbis virus, adenovirus, adeno-associated virus, herpes viruses, SV 40, vaccinia and other DNA viruses. Replication-defective murine retroviral vectors are the most widely utilized gene transfer vectors. Murine leukemia retroviruses are composed of a single strand RNA complexed with a nuclear core protein and polymerase (pol) enzymes, encased by a protein core (gag) and surrounded by a glycoprotein envelope (env) that determines host range. The genomic structure of retroviruses include the *gag*, *pol*, and *env* genes enclosed at by the 5' and 3' long terminal repeats (LTR). Retroviral vector systems exploit the fact that a minimal vector containing the 5' and 3' LTRs and the packaging signal are sufficient to allow vector packaging, infection and integration into target cells providing that the viral structural proteins are supplied in trans in the packaging cell line. Fundamental advantages of retroviral vectors for gene transfer include efficient infection and gene expression in most cell types, precise single copy vector integration into target cell chromosomal DNA, and ease of manipulation of the retroviral genome.

The adenovirus is composed of linear, double stranded DNA complexed with core proteins and surrounded with capsid proteins. Advances in molecular virology have led to the ability to exploit the biology of these organisms to create vectors capable of transducing novel genetic sequences into target cells *in vivo*. Adenoviral-based vectors will express gene product proteins at high levels. Adenoviral vectors have high efficiencies of infectivity, even with low titers of virus. Additionally, the virus is fully infective as a cell free virion so injection of producer cell lines are not necessary. Another potential advantage to adenoviral vectors is the ability to achieve long term expression of heterologous genes *in vivo*.

Mechanical methods of DNA delivery include fusogenic lipid vesicles such as liposomes or other vesicles for membrane fusion, lipid particles of DNA incorporating cationic lipid such as lipofectin, polylysine-mediated transfer of DNA, direct injection of DNA, such as microinjection of DNA into germ or somatic cells, pneumatically delivered DNA-coated particles, such as the gold particles used in a

"gene gun," and inorganic chemical approaches such as calcium phosphate transfection. Particle-mediated gene transfer methods were first used in transforming plant tissue. With a particle bombardment device, or "gene gun," a motive force is generated to accelerate DNA-coated high density particles (such as gold or tungsten) to a high velocity that allows penetration of the target organs, tissues or cells. Particle bombardment can be used in *in vitro* systems, or with *ex vivo* or *in vivo* techniques to introduce DNA into cells, tissues or organs. Another method, ligand-mediated gene therapy, involves complexing the DNA with specific ligands to form ligand-DNA conjugates, to direct the DNA to a specific cell or tissue.

It has been found that injecting plasmid DNA into muscle cells yields high percentage of the cells which are transfected and have sustained expression of marker genes. The DNA of the plasmid may or may not integrate into the genome of the cells. Non-integration of the transfected DNA would allow the transfection and expression of gene product proteins in terminally differentiated, non-proliferative tissues for a prolonged period of time without fear of mutational insertions, deletions, or alterations in the cellular or mitochondrial genome. Long-term, but not necessarily permanent, transfer of therapeutic genes into specific cells may provide treatments for genetic diseases or for prophylactic use. The DNA could be reinjected periodically to maintain the gene product level without mutations occurring in the genomes of the recipient cells. Non-integration of exogenous DNAs may allow for the presence of several different exogenous DNA constructs within one cell with all of the constructs expressing various gene products.

Electroporation for gene transfer uses an electrical current to make cells or tissues susceptible to electroporation-mediated mediated gene transfer. A brief electric impulse with a given field strength is used to increase the permeability of a membrane in such a way that DNA molecules can penetrate into the cells. This technique can be used in *in vitro* systems, or with *ex vivo* or *in vivo* techniques to introduce DNA into cells, tissues or organs.

Carrier mediated gene transfer *in vivo* can be used to transfect foreign DNA into cells. The carrier-DNA complex can be conveniently introduced into body fluids or the bloodstream and then site-specifically directed to the target organ or tissue in the body. Both liposomes and polycations, such as polylysine, lipofectins
5 or cytofectins, can be used. Liposomes can be developed which are cell specific or organ specific and thus the foreign DNA carried by the liposome will be taken up by target cells. Injection of immunoliposomes that are targeted to a specific receptor on certain cells can be used as a convenient method of inserting the DNA into the cells bearing the receptor. Another carrier system that has been used is the
10 asialoglycoprotein/polylysine conjugate system for carrying DNA to hepatocytes for *in vivo* gene transfer.

The transfected DNA may also be complexed with other kinds of carriers so that the DNA is carried to the recipient cell and then resides in the cytoplasm or in the nucleoplasm. DNA can be coupled to carrier nuclear proteins in specifically
15 engineered vesicle complexes and carried directly into the nucleus.

Gene regulation of EM 1 may be accomplished by administering compounds that bind to the EM 1 gene, or control regions associated with the EM 1 gene, or its corresponding RNA transcript to modify the rate of transcription or translation. Additionally, cells transfected with a DNA sequence encoding EM 1 may be
20 administered to a patient to provide an *in vivo* source of EM 1. For example, cells may be transfected with a vector containing a nucleic acid sequence encoding EM 1. The transfected cells may be cells derived from the patient's normal tissue, the patient's diseased tissue, or may be non-patient cells.

For example, tumor cells removed from a patient can be transfected with a
25 vector capable of expressing the EM 1 protein of the present invention, and re-introduced into the patient. The transfected tumor cells produce EM 1 levels in the patient that inhibit the growth of the tumor. Patients may be human or non-human animals. Cells may also be transfected by non-vector, or physical or chemical methods known in the art such as electroporation, ionoporation, or via a "gene gun."

Additionally, EM 1 DNA may be directly injected, without the aid of a carrier, into a patient. In particular, EM 1 DNA may be injected into skin, muscle or blood.

The gene therapy protocol for transfecting EM 1 into a patient may either be through integration of the EM 1 DNA into the genome of the cells, into
5 minichromosomes or as a separate replicating or non-replicating DNA construct in the cytoplasm or nucleoplasm of the cell. EM 1 expression may continue for a long-period of time or may be reinjected periodically to maintain a desired level of the EM 1 protein in the cell, the tissue or organ or a determined blood level.

In addition, the invention encompasses antibodies and antisera, which can be
10 used for testing of novel anti-angiogenic proteins, and can also be used in diagnosis, prognosis, or treatment of diseases and conditions characterized by, or associated with, angiogenic activity or lack thereof. Such antibodies and antisera can also be used to up-regulate angiogenesis where desired, *e.g.*, in post-infarct heart tissue, antibodies or antisera to the EM 1 protein can be used to block localized, native anti-
15 angiogenic proteins and processes, and increase formation of new blood vessels and inhibit atrophy of heart tissue.

Such antibodies and antisera can be combined with pharmaceutically-acceptable compositions and carriers to form diagnostic, prognostic or therapeutic compositions. The term "antibody" or "antibody molecule" refers to a population of
20 immunoglobulin molecules and/or immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antibody combining site or paratope.

Passive antibody therapy using antibodies that specifically bind EM 1 can be employed to modulate angiogenic-dependent processes such as reproduction,
25 development, and wound healing and tissue repair. In addition, antisera directed to the Fab regions of EM 1 antibodies can be administered to block the ability of endogenous EM 1 antisera to bind EM 1.

The EM 1 of the present invention also can be used to generate antibodies that are specific for the inhibitor and its receptor. The antibodies can be either
30 polyclonal antibodies or monoclonal antibodies. These antibodies that specifically

bind to the EM 1 or EM 1 receptors can be used in diagnostic methods and kits that are well known to those of ordinary skill in the art to detect or quantify the EM 1 or EM 1 receptors in a body fluid or tissue. Results from these tests can be used to diagnose or predict the occurrence or recurrence of a cancer and other angiogenic mediated diseases.

The invention also includes use of EM 1, antibodies to EM 1, and compositions comprising EM 1 and/or its antibodies in diagnosis or prognosis of diseases characterized by angiogenic activity. As used herein, the term "prognostic method" means a method that enables a prediction regarding the progression of a disease of a human or animal diagnosed with the disease, in particular, an angiogenesis dependent disease. The term "diagnostic method" as used herein means a method that enables a determination of the presence or type of angiogenesis-dependent disease in or on a human or animal.

The EM 1 can be used in a diagnostic method and kit to detect and quantify antibodies capable of binding EM 1. These kits would permit detection of circulating EM 1 antibodies which indicates the spread of micrometastases in the presence of EM 1 secreted by primary tumors in situ. Patients that have such circulating anti-EM 1 antibodies may be more likely to develop multiple tumors and cancers, and may be more likely to have recurrences of cancer after treatments or periods of remission. The Fab fragments of these anti-EM 1 antibodies may be used as antigens to generate anti-EM 1 Fab-fragment antisera which can be used to neutralize anti-EM 1 antibodies. Such a method would reduce the removal of circulating EM 1 by anti-EM 1 antibodies, thereby effectively elevating circulating EM 1 levels.

The present invention also includes isolation of receptors specific for EM 1. Protein fragments that possess high affinity binding to tissues can be used to isolate the EM 1 receptor on affinity columns. Isolation and purification of the EM 1 receptor is a fundamental step towards elucidating the mechanism of action of EM 1. Isolation of an EM 1 receptor and identification of EM 1 agonists and antagonists will facilitate development of drugs to modulate the activity of the EM 1 receptor,

the final pathway to biological activity. Isolation of the receptor enables the construction of nucleotide probes to monitor the location and synthesis of the receptor, using *in situ* and solution hybridization technology. Further, the gene for the EM 1 receptor can be isolated, incorporated into an expression vector and
5 transfected into cells, such as patient tumor cells to increase the ability of a cell type, tissue or tumor to bind EM 1 and inhibit local angiogenesis.

EM 1 proteins are employed to develop affinity columns for isolation of the EM 1 receptor from cultured tumor cells. Isolation and purification of the EM 1 receptor is followed by amino acid sequencing. Using this information the gene or
10 genes coding for the EM 1 receptor can be identified and isolated. Next, cloned nucleic acid sequences are developed for insertion into vectors capable of expressing the receptor. These techniques are well known to those skilled in the art. Transfection of the nucleic acid sequence(s) coding for EM 1 receptor into tumor cells, and expression of the receptor by the transfected tumor cells enhances the
15 responsiveness of these cells to endogenous or exogenous EM 1 and thereby decreasing the rate of metastatic growth.

Angiogenesis-inhibiting proteins of the present invention can be synthesized in a standard microchemical facility and purity checked with HPLC and mass spectrophotometry. Methods of protein synthesis, HPLC purification and mass
20 spectrophotometry are commonly known to those skilled in these arts. EM 1 proteins and EM 1 receptors proteins are also produced in recombinant *E. coli* or yeast expression systems, and purified with column chromatography.

Different protein fragments of the intact EM 1 molecule can be synthesized for use in several applications including, but not limited to the following; as antigens
25 for the development of specific antisera, as agonists and antagonists active at EM 1 binding sites, as proteins to be linked to, or used in combination with, cytotoxic agents for targeted killing of cells that bind EM 1. The amino acid sequences that comprise these proteins are selected on the basis of their position on the exterior regions of the molecule and are accessible for binding to antisera. The amino and

carboxyl termini of EM 1, as well as the mid-region of the molecule are represented separately among the fragments to be synthesized.

The synthetic protein fragments of EM 1 have a variety of uses. The protein that binds to the EM 1 receptor with high specificity and avidity is radiolabeled and
5 employed for visualization and quantitation of binding sites using autoradiographic and membrane binding techniques. This application provides important diagnostic and research tools. Knowledge of the binding properties of the EM 1 receptor facilitates investigation of the transduction mechanisms linked to the receptor.

EM 1 and EM 1-derived proteins can be coupled to other molecules using
10 standard methods. The amino and carboxyl termini of EM 1 both contain tyrosine and lysine residues and are isotopically and nonisotopically labeled with many techniques, for example radiolabeling using conventional techniques (tyrosine residues-chloramine T, iodogen, lactoperoxidase; lysine residues-Bolton-Hunter reagent). These coupling techniques are well known to those skilled in the art.
15 Alternatively, tyrosine or lysine is added to fragments that do not have these residues to facilitate labeling of reactive amino and hydroxyl groups on the protein. The coupling technique is chosen on the basis of the functional groups available on the amino acids including, but not limited to amino, sulfhydryl, carboxyl, amide, phenol, and imidazole. Various reagents used to effect these couplings include among
20 others, glutaraldehyde, diazotized benzidine, carbodiimide, and p-benzoquinone.

EM 1 proteins are chemically coupled to isotopes, enzymes, carrier proteins, cytotoxic agents, fluorescent molecules, chemiluminescent, bioluminescent and other compounds for a variety of applications. The efficiency of the coupling reaction is determined using different techniques appropriate for the specific
25 reaction. For example, radiolabeling of an EM 1 protein with ^{125}I is accomplished using chloramine T and Na^{125}I of high specific activity. The reaction is terminated with sodium metabisulfite and the mixture is desalted on disposable columns. The labeled protein is eluted from the column and fractions are collected. Aliquots are removed from each fraction and radioactivity measured in a gamma counter. In this
30 manner, the unreacted Na^{125}I is separated from the labeled EM 1 protein. The

protein fractions with the highest specific radioactivity are stored for subsequent use such as analysis of the ability to bind to EM 1 antisera.

In addition, labeling EM 1 proteins with short lived isotopes enables visualization of receptor binding sites *in vivo* using positron emission tomography or
5 other modern radiographic techniques to locate tumors with EM 1 binding sites.

Systematic substitution of amino acids within these synthesized proteins yields high affinity protein agonists and antagonists to the EM 1 receptor that enhance or diminish EM 1 binding to its receptor. Such agonists are used to suppress the growth of micrometastases, thereby limiting the spread of cancer.
10 Antagonists to EM 1 are applied in situations of inadequate vascularization, to block the inhibitory effects of EM 1 and promote angiogenesis. For example, this treatment may have therapeutic effects to promote wound healing in diabetics.

The EM 1 protein of the present invention can also be used as a nutritional source or supplement. Such uses include without limitation use as a protein or amino
15 acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases, the EM 1 protein of the invention can be added to the food of a particular organism, or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be
20 added to the medium in or on which the microorganism is cultured.

The invention is further illustrated by the following examples, which are not meant to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof, which, after reading the
25 description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention and/or the scope of the appended claims.

EXAMPLES

Example 1: Cells and Cell lines

Cell line 786-0 (ATCC No. CRL-1932), a renal clear cell carcinoma line; C-PAE (ATCC No. CCL-209), a bovine pulmonary arterial endothelial cell line and
5 ECV304 (ATCC No. CRL-1998), a human endothelial cell line were all obtained from ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia, 20110-2209, USA). The cell lines were maintained in either DMEM (786-0 and C-PAE) or M199 (ECV304), supplemented with 10% fetal calf serum, 100 U/ml of penicillin, 100 µg/ml of streptomycin and 2 mM L-glutamine.
10 The cDNA clone for mouse endostatin pBACPak 8 was kindly provided by B. R. Olsen, Department of Cellular Biology, Harvard Medical School, Boston. The prokaryotic expression vector pET17b was purchased from Novagen (Madison, Wisconsin, USA). The yeast expression system, *Pichia pastoris* (pPICZαA) was purchased from InVitrogen (San Diego, California, USA). Restriction enzymes and
15 Vent DNA polymerase were purchased from New England Biolabs (Beverly, Massachusetts, USA).

Example 2: Cloning and expression of mouse endostatin and mutants into a prokaryotic system

The gene encoding mouse endostatin was amplified from the pBACPak 8
20 plasmid and expressed initially in the pET expression system. The sequence encoding the carboxy terminal portion of mouse collagen XVIII was amplified by amplification using Vent DNA polymerase, with the endostatin pBACPak 8 vector as a template. The primers used were 5'-GGC ATA TGC ATA CTC ATC AGG ACT TT-3' (SEQ ID NO:3) and 5'-AAC TCG AGA TTT GGA GAA AGA GGT-3'.
25 Amplification was carried out for 30 cycles with the following parameters: 94°C for denaturation, 60°C for annealing, and 72°C for extension, each for 1 minute. The amplified DNA fragment (555 bp) was purified using a QIAquick purification kit, digested with *Nde*I and *Xho*I (these restriction sites are underlined in the primers above), and ligated into the expression vector pET17bhis (Dhanabal *et al.* (1995) *J.*

Immunol. Methods. 182:165-175). Initial transformation was carried out with the host strain HMS174 (Novagen, Madison, Wisconsin, USA). Positive clones were sequenced on both strands. The desired clones were finally transformed into BL21(DE3) (Novagen, Madison, Wisconsin, USA) for expression. The expression of recombinant protein in the pET system was carried out as recommended by the manufacturer (Novagen, Madison, Wisconsin, USA).

A Ni-NTA agarose column was used to purify the recombinant protein. Protein present in inclusion bodies was solubilized in 8 M urea and purified under denaturing conditions as described by O'Reilly *et al.* (1997) (*Cell* 88:277-285). The results are shown in Fig. 1, which is a graph showing the absorbance at 280 nm of the eluted fractions (○). Also plotted is the pH of the elution buffer (●). Fig. 1 shows a small peak around fractions 7-8, a sharp peak around fractions 21-22, and another small peak around fraction 35.

SDS-PAGE analysis of 10-ml samples of selected fractions showed a discrete band at 22-24 kDa under non-reducing conditions. Results are shown in Fig. 2, which shows a 22-24 kDa band for fractions 7 and 8 (lanes 3 and 4, respectively), and also for fractions 21 and 22 (lanes 5 and 6, respectively). In addition, higher molecular weight complexes of 46 and 69 kDa were also observed, which upon reduction with DTT resulted in a discrete band at 22-24 kDa (lane 7 of Fig. 2). The peaks at different pH elutions (pH 4.2 and 3.0) were pooled and dialyzed against decreasing concentrations of urea, and final dialysis was performed in PBS buffer (pH 7.4), at which time most of the proteins precipitated out of solution. Since non-refolded precipitated protein expressed from a similar system had shown biological activity *in vivo*, the exact procedure for "protein refolding" was done as described by O'Reilly *et al.* (1997) (*Cell* 88:277-285). The precipitated protein was used in suspension form for *in vivo* experiments only, with the concentration of protein measured by BCA assay (Pierce Chemical Co., Rockford, Illinois, USA) (solubilized in urea with a suitable blank) and stored at -70°C in small aliquots. Since mouse and human endostatin are conserved at the C-terminus, two small deletions were made. Primers were designed such that either 9 or 17 amino acids

were deleted from the C-terminus of endostatin, resulting in two mutants, designated EM 1 and EM 2, respectively. For EM 1, all 4 of the cysteine residues were left intact. For EM 2, the most C-terminal cysteine was also deleted. The upstream primer for the EM 1 mutant was 5'-TTC CAT ATG CAT ACT CAT CAG GAC
5 TTT CAG GCA-3' (SEQ ID NO:7), and the downstream primer was 5'-TTA GCG
GCC GCC TAC TCA ATG CAG AGG ACG ATG TA-3' (SEQ ID NO:8). The
upstream primer for the EM 2 mutant was 5'-TTC CAT ATG CAT ACT CAT CAG
GAC TTT CAG CCA-3' (SEQ ID NO:9), and the downstream primer was 5'-TTA
GCG GCC GCC TAG TTG TGG CAG CTC GCA GCT TTC TG-3' (SEQ ID
10 NO:10).

The amplified DNA fragments (528 bp for EM 1, 504 bp for EM 2) were purified, digested with *NdeI* and *NotI*, and ligated into a predigested pET28(a) expression vector. The rest of the protocol was carried out as described above. Induction conditions and processing of the bacterial pellet were as described by
15 O'Reilly *et al.* (1997) (*Cell* 88:277-285). The purification of recombinant protein was performed using a Ni-NTA column in the presence of 8 M urea as described in the QIAexpressionist manual (Qiagen, Hilden, Germany). Briefly, the bacterial pellet was solubilized in equilibration buffer (8 M urea, 10 mM Tris and 100 mM sodium phosphate buffer, pH 8.0) for one hour at room temperature. The suspension
20 was sonicated 3-4 times, centrifuged at 10,000x g and the soluble fraction was loaded on a Ni-NTA column pre-equilibrated with the above buffer at a flow rate of 10-20 ml per hour. The column was washed extensively with equilibration buffer. Bound proteins were eluted by lowering the pH of the buffer from 8.0 to 6.3, then to 4.2, and finally to 3.0. For the *in vivo* experiments utilizing endostatin mutants,
25 non-specific proteins binding to the column were removed by an equilibration buffer wash, followed by 10 mM and 25 mM imidazole washes. Bound proteins were eluted in equilibration buffer containing 0.2 M acetic acid. The purified fractions were analyzed by SDS-PAGE and the fractions containing purified endostatin (pH 4.2 and 3.0 for wild type endostatin and equilibration buffer containing 0.2 M acetic
30 acid for endostatin mutants) were pooled and refolded slowly. The final dialysis was

carried out against PBS (pH 7.4) at 4°C. During dialysis the protein precipitated out of solution. It was further concentrated and stored at -70°C in small aliquots. The concentration of protein was determined by the BCA assay (Pierce Chemical Co., Rockford, Illinois, USA).

5 Example 3: Expression of mouse endostatin in *Pichia pastoris*

Pichia pastoris, a methanotropic yeast strain, has many advantages of a higher eukaryotic expression system: (a) the presence of alpha factor signal sequence facilitates secretion of the expressed protein into the medium, (b) the yeast strain (GS115) secretes only very low levels of endogenous host protein which
10 further simplifies the purification process, (c) endotoxin contamination is not an issue, and (d) glycosylation can occur. The pPICZαA vector was selected for expression of mammalian endostatin and its mutants and fragments, because this system produces anti-angiogenic proteins in high titer, and with excellent biological activity, as is described in detail in U.S.S.N. XX/XXX,XXX, "Methods of Producing
15 Anti-Angiogenic Proteins," by Vikas P. Sukhatme, filed December 8, 1998, the entire teachings of which are incorporated herein by reference. When this expression system was used, mammalian endostatin was found to be expressed as a soluble protein (20 kDa) with a peak level of expression noted on the second day after induction.

20 The sequence encoding mouse endostatin was further modified by amplification using Vent DNA polymerase on a template of pET17bhis construct, which contained the mouse endostatin described above. The upstream primer used was 5'-GGG AAT TCC ATA CTC ATC AGG ACT TT-3' (SEQ ID NO:11), and the downstream primer was 5'-AAG CGG CCG CCT ATT TGG AGA AAG AGG T-3'
25 (SEQ ID NO:12). The amplified fragment containing *EcoRI* and *NotI* restriction sites was subcloned into a predigested yeast expression vector. The pPICZαA vector carries an alpha factor secretion signal sequence with a Zeocin marker for antibiotic selection. Initial transformation was done in the 'Top 10' host strain (Invitrogen, San Diego, California, USA). The resultant clones were screened for

the presence of an insert and positive clones were sequenced. The plasmid was then linearized with *SacI* and used for homologous recombination into the yeast host strain GS115 (InVitrogen, San Diego, California, USA). The transformation was carried out by the lithium chloride method as described in the *Pichia* expression manual. Recombinants were selected by plating on YPD plates containing 100 µmg/ml of Zeocin. Clones which grew on YPD/Zeocin plate were tested for expression.

Initial screening was used to identify yeast clones with high levels of expression. The expression of mouse endostatin in large scale was carried out in 2-liter baffled shaker flasks. The overnight culture (A_{600} , 2-6) was used to inoculate 2-liter flasks, with addition of 500 ml of buffered glycerol medium. Cells were grown at 250 rpm at 30°C until A_{600} , 16-20 (2 days). Subsequently, cells were centrifuged at 5000 rpm for 10 minutes, and the yeast resuspended in 300-400 ml of buffered methanol induction medium. The supernatant containing the secreted recombinant protein was harvested on the second, third, and fourth day after induction. After the final harvest, the cell free supernatant was processed immediately.

Example 4: Purification of mouse endostatin via heparin-agarose chromatography

A heparin-agarose column was used for purification, based on data of O'Reilly, *et al.* (1997) (*Cell* 88:277-285). The crude supernatant containing recombinant protein was concentrated by ammonium sulfate precipitation (70%). The precipitated protein was dissolved in 10 mM Tris buffer pH 7.4 containing 150 mM NaCl and dialyzed overnight at 4°C with three changes at 6-8 hour intervals. The dialyzed sample was further concentrated by ultra-filtration using an Amicon concentrator (YM10). A disposable polyprep column (BioRad, Hercules, California, USA) was packed with heparin-agarose resin and equilibrated with 10 mM Tris, 150 mM NaCl, pH 7.4. The concentrated sample was loaded on the column at a flow rate of 20 ml/hour using a peristaltic pump. The column was washed with equilibration buffer until the A_{280} was greater than 0.001. Bound proteins were eluted in 2-ml fractions by a step-wise gradient of NaCl at 0.3 M, 0.6 M, 1 M and 2

M NaCl). The peak fractions from 0.6 M to 1 M were pooled and dialyzed against PBS, pH 7.4. Protein concentration was measured by the BCA assay (Pierce Chemical Co., Rockford, Illinois, USA). The purification process was performed at 4°C in a cold room. Recombinant soluble endostatin expressed from the *Pichia* system was used in all the *in vitro* assays.

Figs. 3 and 4 show the elution profile and SDS-PAGE analysis, respectively, of the purified protein. Fig. 3 shows the fraction number (x-axis), plotted against the absorbance at 280 nm (○) (left y-axis) and against concentration of NaCl (●) (right y-axis) used to elute the fraction. Two distinct peaks were obtained with increasing concentration of NaCl. The first peak at 0.3 M NaCl was small when compared to the major peak at 0.6 M NaCl. Most of the endostatin protein bound to the column as shown by the lack of the protein in the flow-through fraction (Fig. 4, lane 4). The recombinant protein bound tightly and washing with the low salt Tris buffer removed other yeast derived proteins. Protein eluted from the 0.3 M NaCl fraction had a trace amount of endostatin but was contaminated with other host derived high molecular weight protein. The purified protein migrated at 20 kDa which upon reduction migrated at 22 kDa. The protein fractions eluted at 0.6 M and 1 M NaCl were pooled, concentrated and dialyzed against PBS (pH 7.4). The purified protein was further separated by FPLC using a Superose 12 (Pharmacia Biotech, Inc., Piscataway, New Jersey, USA) size separation column. The elution profile from this column showed a single peak. Aliquots of 10 ml from selected fractions were analyzed on a 12% SDS-PAGE gel, and the results are shown in Fig. 4. SDS-PAGE analysis showed the presence of single discrete band of 22-24 kDa corresponding to endostatin. The level of expression was estimated to be in the range of 15-20 mg per liter of culture.

To further characterize the recombinant protein, N-terminal microsequencing was carried out for seven cycles. It showed that the yeast alpha factor signal peptide was processed and cleaved at alanine. The first seven residues (EFHTHQD) of the purified protein after signal peptide cleavage matched exactly the published

sequence of endostatin protein, with the first two residues (EF) derived from linker sequence.

Example 5: Cloning and expression of His.endostatin into the Pichia expression system

5 The coding region of the mouse endostatin construct in the pET17bhis expression vector is preceded by a His.Tag of 10 histidine residues. The coding region, including the His.Tag sequence, was shuttled into pPICZ α A vector via amplification with *Eco*RI and *Not*I sites. Linearization and recombination into the yeast host strain GS115 were done as described above. The cell-free medium was
10 precipitated with 70% ammonium sulfate. Precipitated proteins were dissolved in 50 mM sodium phosphate buffer (pH 8.0) containing 300 mM NaCl and dialyzed in the same buffer at 4°C with three changes at 6-8 hour intervals. A Ni-NTA column was used for purification of the His.endostatin recombinant protein, as described in the QIAexpressionist manual (Qiagen, Hilden, Germany). Bound proteins were eluted
15 with a step-wise gradient of imidazole (10 mM, 25 mM, 50 mM, and 100 mM). The peak fractions from 50 mM and 100 mM imidazole elutions were pooled and dialyzed against PBS buffer, pH 7.4.

 The results are shown in Figs. 5 and 6. Fig. 5 shows the fraction number (x-axis), plotted against the absorbance at 280 nm (○) (left y-axis) and against
20 concentration of imidazole (●) (right y-axis) used to elute the fraction. Several absorbance peaks were observed, the first being the largest, followed by three smaller peaks. The elution profile of His.endostatin from the Ni-NTA column showed that the recombinant protein bound tightly. The yeast-derived host proteins in the culture supernatant did not bind to the column and were removed during the
25 wash. Bound proteins were eluted by a stepwise gradient of imidazole. The non-specifically bound host derived proteins eluted with the addition of 10 mM imidazole (Fig. 5). At 25 mM imidazole, a small fraction of the recombinant protein was eluted along with proteins of higher molecular weight. Final elution with 50 mM and 100 mM imidazole showed a distinct peak. SDS-PAGE analysis is shown

in Fig. 6. The flow-through fraction (lane 3) did not contain any endostatin, indicating that most of the protein bound to the column. Increasing the concentration of imidazole to 10 mM and 25 mM resulted in the elution of non-specific protein. Purified recombinant His.endostatin migrated as a single band
5 corresponding to 22-24 kDa in 50 mM imidazole. A protein with a molecular weight of 22 kDa was seen at 100 mM along with a smaller amount of protein corresponding to 44-46 kDa. The concentration of purified protein was determined by the BCA method. The level of expression was estimated at 15 mg per liter of culture.

10 Example 6: Characterization of recombinant yeast endostatin and polyclonal antibody generation and Western blot analysis

Polyclonal antiserum to mouse recombinant endostatin produced in yeast was raised by immunizing a rabbit with 10 µg of purified protein derived from the *Pichia* expression system. Recombinant endostatin expressed from bacteria and
15 yeast system were separated on a 12% SDS-PAGE gel. The proteins were transferred to PVDF membrane by semi-dry transfer (Trans-blot, BioRad, Hercules, California, USA). The primary antiserum was diluted to 1:4000 in 1x TBS buffer containing 5% non-fat dry milk. Goat anti-rabbit IgG/HRP conjugate was used as a secondary antibody (1:5000). Immunoreactivity was detected by
20 chemiluminescence (Pierce Chemical Co., Rockford, Illinois, USA).

The purified endostatin expressed from the bacterial and yeast expression systems were run under reducing and non-reducing conditions. Fig. 7 shows immunoreactive bands corresponding to endostatin. The size of the protein estimated from the western blot ranges from 22-24 kDa. In addition, the
25 recombinant His.endostatin from yeast and bacteria was probed with a Penta His.monoclonal antibody (Qiagen, Hilden, Germany). The monoclonal antibody showed positive response only with the His.endostatin whereas native endostatin did not show any immunoreactivity. This data confirmed the presence of the His.Tag in the recombinant protein. The antiserum did not show any cross reactivity to human

or mouse angiostatin, demonstrating some degree of immunoreactivity specific to endostatin. Immunoreactivity of the polyclonal antibody was also observed with EM 1 and EM 2 proteins.

Example 7: Endothelial proliferation assay

5 The anti-proliferative effect of endostatin produced in the yeast system was tested using bovine pulmonary artery endothelial cells (C-PAE). Initial experiments were done with different endothelial cell types and various parameters (time of "starvation," serum concentration, concentration and type of mitogenic stimulus (*e.g.*, VEGF vs. bFGF)). C-PAE cells gave the most reproducible response.

10 C-PAE cells were plated in 24-well plates coated with fibronectin (10 µg/ml) at 12,500 cells per well in 0.5 ml DMEM containing 2% FBS. After a 24-hour incubation at 37°C, the medium was replaced with fresh DMEM and 2% FBS containing 3 ng/ml of bFGF (R & D systems, Minneapolis, Minnesota, USA) with or without recombinant mouse endostatin. The cells were pulsed with 1 µCi of
15 ³H-thymidine for 24 hours. Medium was aspirated, cells were washed three times with PBS, and then solubilized by addition of 1.5 N NaOH (100 µl per well) and incubated at 37°C for 30 minutes. Cell-associated radioactivity was determined with a liquid scintillation counter. The experiment was repeated 5 times under identical conditions, with similar results each time.

20 A dose dependent inhibition of bFGF induced proliferation was observed. The results are shown in Fig. 8, which is a graph showing concentration of yeast-derived soluble endostatin (○) and yeast-derived soluble His.endostatin (●) along the x-axis, and incorporation of ³H-thymidine on the y-axis. In general, incorporation decreased steadily with increasing concentration of endostatin. The
25 inhibition range (30-94% of control) was seen with increasing concentrations of endostatin (0.1 µg/ml to 10 µg/ml), with an ED₅₀ value in the range of 600-700 ng/ml. A similar inhibitory effect on C-PAE cells was seen when His.endostatin from yeast was tested in the above assay, as is shown in the graph in Fig. 8. Incorporation of ³H-thymidine dropped steadily with increasing concentration of

either yeast-derived soluble endostatin (○) and yeast-derived soluble His.endostatin (●).

The recombinant protein did not inhibit the proliferation of the renal cell carcinoma cells (786-0 and A498) at concentrations ranging from 0.5 µg/ml to 10 µg/ml, as shown in Fig. 9. Fig. 9 is a bar chart, showing the incorporation of ³H-thymidine in 786-0 cells (open bars), and A498 cells (shaded bars). The recombinant endostatin also did not have an effect on IMR90 and NIH3T3 fibroblasts.

Example 8: Endothelial cell migration assay

Since C-PAE cells do not migrate in response to bFGF and VEGF, ECV304 cells were used with different concentrations of endostatin using bFGF as a stimulus. To determine the ability of recombinant endostatin to block migration of ECV304 cells towards bFGF, a migration assay was performed using 12-well Boyden chemotaxis chambers (Neuro-Probe, Inc., Cabin John, Maryland, USA) with a polycarbonate membrane (25 x 80 mm PVD free, 8 µ pores, Poretics Corp., Livermore, California, USA). The non-specific binding of growth factor to the chambers was prevented by coating the chambers with a solution containing 0.5% gelatin, 1 mM CaCl₂ and 150 mM NaCl at 37°C overnight. ECV304 cells were grown in 10% FBS containing 5 ng/ml DiI (1,1'-dioctadecyl-3, 3, 3', 3'-tetramethylindocarbocyanine perchlorate DiIC18, Molecular Probes, Eugene, Oregon, USA) overnight and washed with PBS containing 0.5% BSA. Following trypsinization, the cells were counted using Coulter-Counter Z1, (Luton, U.K.) and diluted to 300,000 cells/ml in Medium 199 (Life Technologies, Gibco/BRL, Gaithersburg, Maryland, USA) containing 0.5% FBS. The lower chamber was filled with Medium 199 containing 25 ng/ml bFGF. The upper chamber was seeded with 15,000 cells/well with different concentrations of recombinant endostatin. Cells were allowed to migrate for 4 hours at 37°C. At that time, the cells on the upper surface of the membrane were removed with a cell scraper and the (migrated) cells on the lower surface were fixed in 3% formaldehyde and washed in PBS. Images of

the fixed membrane were obtained using fluorescence microscopy at 550 nM with a digital camera and the number of cells on each membrane was determined using the OPTIMAS (version 6.0) software (Media Cybernetics, L.P., Silver Spring, MD, USA).

5 Addition of endostatin resulted in a dose-dependent inhibition of migration, as shown in Figs. 10A and 10B, and Fig. 11. Figs. 10A and 10B are photomicrographs showing inhibition of endothelial cell (ECV304) migration by soluble mouse endostatin using bFGF (25 ng/ml) as a stimulus. Fig. 10A shows migrated endothelial cells in the control (+ bFGF, no endostatin), and Fig. 10B
10 shows migrated endothelial cells treated with endostatin (20 µg/ml) with bFGF.

Fig. 11 is a bar chart showing inhibition of endothelial cell migration with different concentrations of endostatin. Relative cell migration is shown on the y-axis, and treatment (control, 25 ng/ml bFGF, and endostatin at 20, 10, 5, 2.5, and 1 µg/ml) on the x-axis. Each treatment was done in duplicate. In each well, the
15 number of cells migrated was counted in three different areas and the average obtained. Each value is a mean from representative experiments and error bars represent standard deviations. At a concentration less than 1 µg/ml, marginal inhibition of migration was noted, whereas at 10 µg/ml, 60% inhibition of endothelial cell migration was observed. These studies are the first to show
20 endostatin's effect on cell migration. Endostatin's action on migration of two non-endothelial cell lines was also assessed. No effect was seen on inner medullary collecting duct renal cells (IMCD), and some effect (15% at 5 µg/ml and 50% at 20 µg/ml) was noted in the IC-21 macrophage precursor cell line, suggesting that at high concentration, endostatin may block cell migration in some cell types.

25 Example 9: Chorioallantoic membrane (CAM) assay

The ability of mouse endostatin to block bFGF induced angiogenesis *in vivo* was tested using the chorioallantoic membrane (CAM) assay. Fertilized white Leghorn chicken eggs (SPAFAS, Inc., Norwich Connecticut, USA) were opened on 100 mm² petri dishes and allowed to grow until day 11 in a humidified incubator at

38°C. Pellets containing vitrogen (Collagen Biomaterials, Palo Alto, California, USA) at a concentration of 0.73 mg/ml were supplemented with either: vehicle alone; VEGF (250 ng/pellet), VEGF (250 ng/pellet) and endostatin (20 to 0.5 µg/pellet), bFGF (50 ng/pellet), or bFGF (50 ng/pellet) and endostatin (20 to 0.5 µg/pellet). The pellets were allowed to polymerize at 37°C for 2 hours. The pellets were placed on a nylon mesh and oriented on the periphery of the CAM. Embryos were returned to the incubator for 24 hours. Invasion of new capillaries on the collagen mesh was assessed by injection of FITC-dextran into the circulation of the chicken embryo. At the end of the experiment, the meshes were dissected and evaluation of vascular density was done using the program NIH Image v1.59 according to the method of Iruela-Arispe *et al.* (1997) (*Thromb. Haemost.* 78:672-677). Assays were performed in triplicate and four independent experiments were conducted.

Endostatin was able to suppress the angiogenic response mediated by both bFGF and VEGF, as shown in Figs 12 and 13. Fig. 12 is a set of three photomicrographs showing the vascular density for vehicle alone (Fig. 12A), VEGF alone (250 ng/pellet, Fig. 12B), and VEGF and endostatin (Fig. 12C). Figs. 13A and 13B show that the inhibition was dose-dependent. Fig. 13A is a bar chart showing concentration of recombinant protein (x-axis), plotted against angiogenic inhibition in response to VEGF (y-axis). Fig. 13B is a bar chart showing concentration of recombinant protein (x-axis), plotted against angiogenic inhibition in response to bFGF (y-axis). All of the counts were normalized to the negative control. Both charts show a steady increase of inhibition of angiogenesis in response to increasing concentrations of endostatin. Blocking of the VEGF response was somewhat more effective (47%) than suppression of the bFGF response (39%), both at 20 µg/mesh.

Example 10: Neutralization of endostatin's inhibitory effect

The specificity of endostatin's inhibitory effect was demonstrated by neutralization studies using endothelial proliferation and CAM assays. In the endothelial proliferation assay, the endostatin was pre-incubated with an excess of

polyclonal antiserum or purified antibody (IgG) for 1 hour at room temperature and then added to the C-PAE cells in the presence of 3 $\mu\text{g/ml}$ bFGF. Pre-immune serum was used as negative control. In addition, purified IgG and endostatin antibody alone were also used as a control. DNA synthesis was measured by adding 1
5 $\mu\text{Ci/well}$ ^3H -thymidine for 24 hours and the cell-associated radioactivity was measured as described above. For the CAM assay, endostatin (10 μg) and antiserum (50 μg) were pre-incubated overnight end-over-end at 4°C prior to preparation of the pellets. Controls for these experiments included IgG alone and pre-immune serum alone. Evaluation of the angiogenic responses in the two assays were determined as
10 indicated above.

Fig. 14 is a bar chart showing neutralization of the inhibitory effect of mouse endostatin by polyclonal antiserum in the endothelial proliferation assay. Incorporation of ^3H -thymidine is shown on the y-axis, and treatment (control, 10 μg endostatin, 10 μg endostatin + antiserum, 5 μg endostatin, 5 μg endostatin +
15 antiserum, pre-immune serum, endostatin antiserum, and endostatin IgG) on the x-axis). Each value is a mean from triplicate culture and error bars represent standard deviation. Fig. 15 is a pair of photomicrographs showing the results of the CAM assay. Fig. 15A shows the effect of VEGF and endostatin (10 μg), and Fig. 15B shows the effect of VEGF, endostatin (10 μg), and polyclonal antiserum (50 μg).
20 Both Fig. 14 and Fig. 15 demonstrate that the inhibitory effect of endostatin can be suppressed by incubation with specific antiserum. Anti-endostatin antiserum blocked the suppressive effect by 95%. The pre-immune serum and endostatin antibody alone did not have a stimulatory effect, nor did normal rabbit IgG.

Example 11: Inhibition of primary 786-0 RCC tumors in nude mouse model

25 Male nude mice of 6-8 weeks of age were injected subcutaneously in the right flank with 2 million 786-0 cells in a 100 μl volume. Tumors appeared approximately two weeks after implantation. Tumor size was measured using calipers and tumor volume was calculated using the standard formula of:

$$\text{tumor volume} = ab^2 \times 0.52$$

where a = length of the tumor, and b = width of the tumor (O'Reilly *et al.* (1994) *Cell* 79:315-328). The tumor volume ranged from 350 mm³ to 400 mm³. The animals were randomized and each group had five mice with comparable tumor size within and among the groups. Treatment was started with recombinant endostatin (bacterial or yeast versions) with each mouse receiving 10 mg/kg body weight of recombinant protein daily, administered for a period of ten days via intraperitoneal injection. Control animals received PBS each day. Tumor size in all groups was measured on alternate days and tumor volume was calculated. The treatment was terminated on day 10 and animals were sacrificed and tumors from each mouse removed and fixed in 10% buffered formalin.

The results are shown in Figs. 16, 17 and 18. Fig. 16 is a graph showing the inhibition of 786-0 tumor growth by systemic treatment with recombinant endostatin. Time in days after treatment is shown on the x-axis and tumor volume in mm³ is shown on the y-axis. Intraperitoneal injection of endostatin was given at 10 mg/kg/day, starting on day 1 (arrow). Each time point represents the average of five mice in each group and the error bar represents S.E.M. Treatments are control PBS (○), endostatin from yeast (●), His.endostatin from yeast (x) and His.endostatin from bacteria (□). On the fifth day after treatment there was a difference between control (963 mm³) and treated tumors. Yeast endostatin-treated tumors were 405 mm³, bacterially-produced endostatin-treated tumors were 442 mm³, and His.endostatin-treated tumors were 462 mm³.

Figs. 17A through 17E are a set of five photographs of 786-0 tumors treated with recombinant endostatin. Figs. 17A and 17B are control tumors, Fig. 17C shows a tumor treated with yeast-derived endostatin, Fig. 17D shows the effect of His.endostatin from bacteria, and Fig. 17E shows a tumor treated with His.endostatin from yeast. At the end of the treatment period, tumors from control and treated groups were examined grossly under a dissecting microscope. Tumors from the control group were in general larger and more highly vascularized. A 2.5-fold decrease in tumor volume was observed on the fifth day after treatment between control and treated tumors (Figs. 16 and 17). The growth of the tumor was

suppressed in all the treatment groups, and a slower growth rate was seen compared to the control group. Bacterial- (His.Tag) or yeast-derived (with or without His.Tag) endostatin at a dose of 10 $\mu\text{g/kg}$ all worked equally well. On the tenth day after treatment, the tumor volume in the control animals was 1490 mm^3 , whereas in the
5 treated group it was in the range of 480-570 mm^3 (p value < 0.005). Endostatin administration did not inhibit tumor growth completely; the growth of the tumors slowed, with a marginal increase in volume during the treatment period.

Example 12: Two closely related C-terminus endostatin mutants generated in *E. coli* show markedly differing in vivo activity in RCC

10 The RCC tumor model described above was used in a second set of experiments with endostatin and mutants EM 1 and EM 2, produced in the prokaryotic system in Example 1, above. The daily dosage was 20 mg of the protein per kg body weight, injected intraperitoneally. The initial tumor volume was 150-200 mm^3 . Wild type endostatin, also produced in the pET28(a) vector, was
15 given at 20 mg/kg body weight for the experiment as a positive control and PBS was given as a negative control.

The results are Shown in Fig. 18, which is a graph showing days after treatment on the x-axis, and tumor volume on the y-axis. Each time point represents the average of five mice in each group. Treatments were control PBS (○), wild type
20 His.endostatin from bacteria (dotted line, ●), EM 1 from bacteria (dashed line, △), and EM 2 from bacteria (solid line, ◆). Intraperitoneal injection was started on day 1 (arrow). Nine days after treatment, the difference between groups was apparent (Fig. 18). On the eleventh day after treatment, the tumor volume in the control group (397 mm^3) was approximately twice that of the two treated groups: endostatin
25 (182 mm^3) or EM 1 (259 mm^3). However, on the same day, the tumor volume of the EM 2-treated group (389 mm^3) was similar to that of the control group (397 mm^3). Significance was at the 90% confidence level between the EM 2 and endostatin groups and 95% confidence level between endostatin and control groups. Dropping the value of the largest and smallest tumors on day 11 in each group increased the

confidence level to 95% between EM 2 and EM 1 and between EM 2 and endostatin. Therefore, the EM 1 protein retained the native biological activity of endostatin, whereas EM 2, with its further deletion of 8 amino acids, did not. In addition, two of the five mice in the endostatin group and one of the five in the EM 1 group had no detectable tumor at the end of the treatment period.

Example 13: Annexin V - FITC assay

Annexin V, a calcium dependent phospholipid binding protein with a high affinity for phosphatidylserine (PS) was used to detect early stage apoptosis. After initiation of apoptosis, most cell types translocate the membrane phospholipid phosphatidylserine (PS) from the inner surface of the plasma membrane to the outside. PS can be detected by staining with an FITC conjugate of Annexin V, 38 kDa protein that binds naturally to PS. During PCD, PS externalization typically precedes membrane bleb formation and DNA fragmentation.

Briefly, 200,000 cells were plated onto a fibronectin-coated 6-well plate in DMEM containing 2% FBS and 3 ng/ml of bFGF. Different concentrations of recombinant mouse endostatin were added to each well, and cells were harvested and processed 18 hours after treatment. For the time course study, 10 μ g/ml of endostatin was added and cells were processed after 3, 4, 6, 12, and 18 hours. Human recombinant TNF- α (40 ng/ml) was used as a positive control. The cells were washed in PBS and resuspended in binding buffer (10 mM HEPES/NaOH, pH 7.4, 140 mM NaCl, 2.5 mM CaCl_2). Annexin V - FITC was added to a final concentration of 100 ng/ml, and the cells were incubated in the dark for 10 minutes, then washed again in PBS and resuspended in 300 μ l of binding buffer. 10 μ l of propidium iodide (PI) was added to each sample prior to flow cytometric analysis. The cells were analyzed using a Becton Dickinson FACStar plus flow cytometer. Electronic compensation was used to eliminate bleed-through fluorescence. In each sample, a minimum of 10,000 cells were counted and stored in listmode. Data analysis was performed with standard Cell Quest software (Becton-Dickinson). The quadrant settings were set so that the negative control allowed less than 1%

positivity. Endostatin was added to non-endothelial cells (NIH3T3 and 786-0) at 10 $\mu\text{g/ml}$ and the cells were processed and analyzed as described above.

Endostatin at 10 $\mu\text{g/ml}$ showed a distinct shift in Annexin fluorescence intensity. The mean fluorescence intensity difference between control and
5 endostatin treated cell was significant ($p = 0.01$) at 5 and 10 $\mu\text{g/ml}$. The shift in fluorescence intensity was similar for endostatin at 10 $\mu\text{g/ml}$ and the positive control $\text{TNF-}\alpha$ (40 ng/ml). Concentrations of endostatin below 0.1 $\mu\text{g/ml}$ did not show any significant Annexin V positivity. In order to investigate the earliest time point at which endostatin caused externalization of PS, we conducted a time course
10 experiment (Fig 2B). The effect of endostatin was significant ($p = 0.01$) at 12 hours after treatment. Time points before 6 hours did not show a difference between control and treated samples.

Morphological examination of FACS analyzed samples with fluorescence microscopy (Nikon) showed Annexin V staining localized to the cell membrane at
15 12 hours with no staining in the cytoplasm. During this period, the majority of the cells were negative for PI, implicating the early stage of apoptosis. With increased exposure time (24-36 hours), in addition to membrane staining with Annexin V some of the cells turned positive for PI, consistent with a more advanced stage of apoptosis.

20 Similar levels of Annexin V staining were observed in two other endothelial cell lines studied, BAE and BCE. We have also tested the effect of human endostatin on these three bovine endothelial cell lines. We failed to detect Annexin V staining in the presence of human endostatin added to these cells, whereas when human endothelial cell lines were used (HUVE and HMVE-L), it resulted in a
25 marked shift in Annexin V fluorescence (manuscript in preparation). These data indicate that apoptosis, as assessed by Annexin V staining, occurs in diverse endothelial cells in response to mouse and human endostatin.

With regard to non-endothelial cells, 786-0 and NIH3T3 cells failed to show any distinct annexin positivity. In addition, other non-endothelial cells (IMR-90,
30 A10 and H9c2 (2-1)-myoblast) were screened and no effect of endostatin was found.

Based on these results, endostatin's action appears to be selective for endothelial cells.

Example 14: Caspase 3 assay

Caspase 3 (CPP32) is an intracellular protease activated early during
5 apoptosis of mammalian cells and initiates cellular breakdown by degrading specific structural, regulatory, and DNA repair proteins. This protease activity can be measured spectrophotometrically by detection of the chromophore (p-nitroanilide) after cleavage from the labeled substrate (DEVD-pNA).

This assay was performed in either a 75-cm² tissue culture flask or in
10 fibronectin-coated 6-well plates. The 6-well plates were seeded with $0.5-1 \times 10^6$ cells per well, and the flasks were seeded with 2×10^6 cells. The cells were maintained overnight in DMEM with 10% FBS. The following day, the old medium was replaced with fresh medium (2% FBS), and the cells were incubated overnight at 37°C. Following starvation, the cells were stimulated with bFGF (3 ng/ml) in
15 DMEM (2% FBS). Along with bFGF, yeast endostatin (10 µg/ml final concentration) was added and the cells grown for 24 hours. For the control plate, only the PBS buffer was added. As a positive control, TNF-α was used at a final concentration of 20 ng/ml. After 24 hours, the supernatant cells were centrifuged and collected. The wells (flasks) were trypsinized to collect the attached cells and
20 combined with the supernatant cells. The cells were counted and resuspended in cell lysis buffer (Clontech, Palo Alto, California, USA) at a concentration of 4×10^7 cells/ml. The rest of the protocol followed the manufacturer's instruction (Clontech, Palo Alto, California, USA). A specific inhibitor of caspase 3, DEVD-fmk, was used to confirm the specificity as suggested by the manufacturer. The absorbance
25 was measured in a microplate reader (BioRad, Hercules, California, USA) at 405 nm. Fold-increase in protease activity (caspase 3) was determined by comparing the results of the induced sample (yeast endostatin or TNF-α) with the uninduced control. Similarly non-endothelial cells (NIH3T3 and H9c2 (2-1)-myoblast were used and analyzed as described above.

A time course experiment was first performed with 10 $\mu\text{g/ml}$ of endostatin, looking for an increase in caspase 3 activity. There was no difference in caspase 3 activity between the treated and the control samples at 2, 4, 8, and 14 hours. However, caspase 3 activity 24 hours after treatment with endostatin was elevated
5 over controls. The caspase activity of the endostatin and $\text{TNF-}\alpha$ (positive control) treated samples is shown in Fig. 21. When compared to controls, endostatin treated cells showed a 1.8-fold increase in caspase 3 activity after 24 hours, whereas $\text{TNF-}\alpha$ gave a comparable (1.75-fold) increase. The assay was repeated at least five times with similar results. When a specific inhibitor of caspase 3 (DEVD-fmk) was
10 included in the same samples, the protease activity was at baseline (comparison of the dark box to the corresponding white box), indicating that the increase in the measured activity was specific for caspase 3. Fig. 22 shows that for NIH3T3 cells only, a marginal increase in caspase-3 was seen, whereas in myoblast cells there was no difference in caspase-3 levels between treated and control cultures.

15 Example 15: Microscopic detection of TUNEL staining

Fragmentation of nuclear DNA is one of the distinct morphological changes occurring in the nucleus of an apoptotic cell. A TUNEL (terminal deoxynucleotidyl transferase-mediated dUTP nick-end-labeling) assay was performed on endostatin, $\text{TNF-}\alpha$ treated and control cells. For adherent cells, C-PAE cells were seeded at a
20 density of 5,000 cells per well on fibronectin coated (10 $\mu\text{g/ml}$) Lab-Tek chamber slides and grown in 0.4 ml of DMEM medium with 10% FBS. After two days, the old medium was aspirated and fresh DMEM with 2% FBS was added and the cells were starved overnight. The following day, 0.36 ml of new medium (with 2% FBS) containing 3 ng/ml bFGF was added along with yeast endostatin (10 $\mu\text{g/ml}$) or
25 $\text{TNF-}\alpha$ (20 ng/ml). For control samples, fresh medium (2% FBS) containing bFGF (3 ng/ml) was added. Following induction (24 hours), the slides were washed twice with PBS, and subsequently fixed in fresh 4% formaldehyde/PBS at 4°C for 25 minutes. The slides were washed in PBS and the cells permeabilized in 0.2% Triton

X-100/PBS for 5 minutes on ice, then washed with fresh PBS twice for 5 minutes each at room temperature, and the TUNEL assay performed as described below.

The TUNEL assay was performed as described in the ApoAlert DNA fragmentation assay kit user manual (Clontech, Palo Alto, California, USA), except
5 that the final concentration of propidium iodide (Sigma, St. Louis, Missouri, USA) used was 1 $\mu\text{g/ml}$. After the assay, a drop of anti-fade solution was added and the treated portion of the slide was covered with a glass coverslip with the edges sealed with clear nail polish. Slides were viewed immediately under a fluorescent
10 microscope using a dual filter set for green (520 nm) and red fluorescence (>620 nm). The images were captured using a digital microscope (Nikon Microphot-SA) and processed using SPOT software version 1.1.02. For the positive control (TNF- α), 5 fields random were chosen, and for the samples, 15 random fields were chosen. The number of green and red cells per field were then counted, and the percent of green divided by the number of red cells in a given field was determined.
15 An average (with S.E.M.) of the different fields was then calculated.

For cells in suspension, floating cells were collected by centrifugation at 300x g for 10 minutes at 4°C. The old medium was aspirated, and the cells were resuspended in 500 ml of PBS (pH 7.4). Cells were centrifuged again, the PBS removed, and the pellet was resuspended in 75 ml of fresh PBS. Resuspended cells
20 were spread on a poly-L-lysine coated slide (Jersey lab supply) using a clean slide. The cells were fixed by immersing the slides in fresh 4% formaldehyde/PBS at 4°C for 25 min. The rest of the protocol was carried out as described above.

In the presence of the enzyme TdT, both endostatin and TNF- α treated slides showed numerous positive cells under green fluorescence, whereas no positive cells
25 were seen in the control. Without the enzyme, the endostatin treated slide showed background cell fluorescence.

The number of apoptotic cells in several fields were counted, and the percent of apoptotic cells (green divided by the number of red cells per field) is plotted in Fig. 23, which is a bar graph. The apoptosis rate in the control cells was 1.24%. In
30 the endostatin treated cells, a 30-fold increase in the apoptosis rate was observed in

suspension cells (38.3%), while a 15-fold increase was observed in the attached cells (19.4%). With TNF- α , the apoptosis rate was 6.4%. In contrast, the percent of TUNEL-positive in the angiostatin treated BCE (bovine adrenal cortex capillary endothelial) cells was 2% when compared with the control cells (1.2%), a 1.6 fold increase, suggesting that endostatin is a stronger apoptotic agent than angiostatin.

Example 16: Bcl-2 and Bax expression by western blot analysis

C-PAE cells (1×10^6) were seeded in 10 cm petri dishes precoated with fibronectin (10 μ g/ml) in the presence of 2% FBS containing 3 ng/ml bFGF. Endostatin was added at 10 μ g/ml, and cells were harvested at 12, 24, and 28 hours after treatment. Cells were washed thrice in PBS buffer pH, 7.4 and the cells were resuspended in 1 ml of 1x EBC buffer (50 mM Tris-HCl, pH 8.0, 120 mM NaCl, 1% Nonidet P-40) containing freshly added complete protease inhibitor tablet (Boehringer Mannheim), 100 mg/ml Pefabloc, 1 mg/ml Pepstatin. The protein concentration in whole cell lysate was measured by the BCA method. 30 mg of whole cell extract was loaded onto a 4-15% gradient polyacrylamide gel. Transfer was performed using a semi-dry transblot apparatus (BioRad, Hercules, California, USA). The membrane was blocked in wash buffer (1x TBS) with 5% non fat dry milk and incubated at 37°C for 1 hour. Goat antibody directed against human Bcl-2 (N-19) {sc-492-G} was purchased from Santa Cruz Biotechnology (Santa Cruz, California, USA). Affinity purified mouse polyclonal antibody against Bax (B-9) {sc7490} and Bcl-XS/L {sc1690} were purchased from the same manufacturer. Polyclonal anti-actin antibody (Sigma, St. Louis, Missouri, USA) was used to normalize for protein loading. Secondary antibodies were anti-goat, mouse and rabbit immunoglobulin conjugated to horseradish peroxidase (Amersham Corp., Arlington Heights, Illinois, USA). The immunoreactivity was detected with an enhanced chemiluminescence reagent (Pierce Chemical Co., Rockford, Illinois, USA). Images were scanned using a flat bed scanner (Scan-Jet 4C) and quantitated by the NIH Image 1.61 software. Normalization was done by dividing the Bcl-2 signal by that of actin within each experiment.

Anti-apoptotic members such as Bcl-2 and Bcl-X_L prevent PCD in response to numerous stimuli. Conversely, pro-apoptotic proteins such as Bax and Bak can accelerate cell death; and in certain cases, they are sufficient to cause apoptosis independent of additional signals. Whole cell extract of endostatin treated and control C-PAE cells were tested for Bcl-2 and Bax expression levels. In growth arrested C-PAE cells, Bcl-2 expression was high. It was relatively constant up to 28 hours; in contrast, endostatin treated cells showed marked decrease in Bcl-2, as is show in Fig. 24A. Densitometry revealed that the levels of Bcl-2 compared to control was 1.2, 1.5, and 3 fold less at 12, 24, and 28 hours respectively after treatment, with actin levels used as normalization controls. In contrast, Bax expression was similar between control and treated cultures (Fig. 24B).

Bcl-2 protein was not detected in both NIH3T3 and IMR-90 cells. Bax expression levels were not affected by endostatin treatment in these cell lines, as is shown in Figs. 25A and 25B. In C-PAE cells, at the early time point (12 hours), Bcl-XL level was reduced by 2 fold, whereas in NIH3T3 cell its expression was unchanged (Figs. 25C and 25D). Interestingly, we have detected only the larger pro-apoptotic form of Bcl-X in C-PAE whereas, in NIH3T3 both smaller and larger forms were detected.

These findings suggest that endostatin exerts its regulatory activity by altering Bcl-2 expression. Interestingly, VEGF has been shown to augment Bcl-2 levels in endothelial cells. Since endostatin antagonizes VEGF's proliferative effects, Bcl-2 appears to be one point of regulation. Recent studies indicate that the Bcl-2 protein binds to other proteins, such as Bax, Bcl XS, Bik and Bad, which ultimately enhance cell survival (Newton, K, and Strasser, A. (1998) *Curr. Opin. Genet. Dev.* 8:68-75; Jacobson, M.D. (1997) *Curr. Biol.* 7:R277-81). The function of another Bcl-2 homologue, Bax remains enigmatic. Bcl-2 and Bcl-X_L function through heterodimerization with Bax, and overexpression of Bax accelerates apoptosis. Recently, it was shown that FGF-2 inhibited endothelial cell apoptosis by Bcl-2 dependent and independent mechanisms. In our study, we did see differences in Bcl-2 (and Bcl-X_L) expression in endostatin treated cultures but no difference in

Bax levels. It is possible that Bcl-2 may act independently of Bax, as has been shown for T cells.

EQUIVALENTS

While this invention has been particularly shown and described with
5 references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

CLAIMS

What is claimed is:

1. An isolated anti-angiogenic peptide, wherein the C-terminal of the isolated peptide comprises the amino acid sequence SYIVLCIE.
- 5 2. Isolated EM 1, comprising a mutated endostatin protein, wherein the comprises a deletion of nine consecutive amino acids from the C-terminus of the endostatin protein, and wherein the isolated EM 1 is characterized as having anti-angiogenic activity.
- 10 3. The isolated EM1 of Claim 2, wherein the C-terminus of the isolated EM 1 comprises the amino acid sequence SYIVLCIE.
4. The isolated EM 1 of Claim 2, wherein the deletion of nine consecutive amino acids comprises the amino acid sequence NSFMTSFSK.
- 15 5. The isolated polynucleotide of Claim 1, comprising;
 - (a) the nucleotide sequence of SEQ ID NO.: ;
 - (b) a sequence complementary to the nucleotide sequence of SEQ ID NO.: ; and
 - (c) a sequence that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO.: .
- 20 6. An isolated polynucleotide, comprising the nucleotide sequence amplified by the primers of SEQ ID NO:7 and SEQ ID NO:8.
7. An isolated polynucleotide of Claim 3, wherein the polynucleotide is operably linked to an expression control sequence.

8. A host cell transformed with the polynucleotide of Claim 7.
9. The host cell of Claim 8, where the cell is selected from the group comprising bacterial, yeast, mammalian, insect or plant cells.
10. A process for producing a protein encoded by the polynucleotide of Claim 3,
5 wherein the process comprises:
 - (a) growing a culture of a host cell transformed with the polynucleotide of Claim 3, where the host cell is selected from the group comprising bacterial, yeast, mammalian, insect or plant cells; and
 - (b) purifying the protein from the culture;10 thereby producing the protein encoded by the polynucleotide of Claim 3.
11. A fusion protein, comprising two or more protein molecules, and further comprising the EM 1 of Claim 3.
12. The fusion protein of Claim 11, further comprising at least one protein
15 molecule selected from the group comprising: restin, endostatin, angiostatin, apomigren, or EM 1.
13. A composition comprising, as a biologically active ingredient, the EM 1 of Claim 3.
14. The composition of Claim 23, and a pharmaceutically-compatible carrier.
15. A composition comprising, as a biologically active ingredient, the fusion
20 protein of Claim 11.
16. A composition comprising, as a biologically active ingredient, the fusion protein of Claim 12.

17. A method for inhibiting angiogenic activity in mammalian tissue, the method comprising contacting the tissue with a composition comprising the EM 1 of Claim 3.
 18. A method of using the composition of Claim 17 to treat a disease, the method comprising administration of the composition to a patient with a disease characterized by angiogenic activity.
 19. The method of Claim 18, wherein the disease is selected from the group comprising angiogenesis-dependent cancers, benign tumors, rheumatoid arthritis, psoriasis, ocular angiogenesis diseases, Osler-Webber Syndrome, myocardial angiogenesis, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma, wound granulation, intestinal adhesions, atherosclerosis, scleroderma, hypertrophic scars, cat scratch disease, *Helicobacter pylori* ulcers, dialysis graft vascular access stenosis, contraception, and obesity.
 20. The method of Claim 19, wherein the disease is cancer.
 21. The method of Claim 20, wherein the disease is renal cancer.
 22. A method of using a composition comprising the isolated EM 1 of Claim 3 to induce apoptosis in a cell or tissue, comprising contacting the cell or tissue with the composition.
 23. A method of using any of the compositions of Claims 13 to 16 to treat a disease, the method comprising administration of the composition to a patient with a disease characterized by angiogenic activity.
 24. The method of Claim 23, wherein the disease is cancer.
-

25. The method of Claim 24, wherein the disease is renal cancer.
26. A process for providing a mammal with EM 1 protein, the process comprising introducing mammalian cells into a human, said mammalian cells having been treated *in vitro* to insert therein the polynucleotide encoding the amino acid sequence comprising EM 1 and expressing *in vivo* in said mammal a therapeutically effective amount of the EM 1 protein.
27. The process of Claim 26, wherein the cells are lymphocytes.
28. The process of Claim 27, wherein the lymphocytes are chosen from the group comprising T-lymphocytes and B-lymphocytes.
29. The process of Claim 26, wherein the cells are chosen from the group comprising: blood cells, TIL cells, bone marrow cells, vascular cells, tumor cells, liver cells, muscle cells, fibroblast cells.
30. The process of Claim 26, wherein the polynucleotide is inserted into the cells by a viral vector.
31. A process for producing an isolated polynucleotide, the process comprising the steps of:
- (a) preparing one or more polynucleotide probes that hybridize under conditions of moderate stringency to a nucleotide sequence selected from the group consisting of:
 - (i) SEQ ID NO:1, from nucleotide 1 to nucleotide 525;
 - (ii) an isolated polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2, from amino acid 1 to amino acid 175; and
 - (b) hybridizing said probe(s) to mammalian DNA; and

(c) isolating the DNA polynucleotide detected with the probe(s);
wherein the nucleotide sequence of the isolated polynucleotide corresponds
to the nucleotide sequence of SEQ ID NO:1, from nucleotide 1 to nucleotide
525.

- 5 32. An isolated polynucleotide produced according to the process of Claim 31.
33. An isolated polynucleotide comprising the polynucleotide of Claim 32.
34. Antibodies to the isolated anti-angiogenic mutant fragment of endostatin of
Claim 2.
35. An isolated mutant, derivative, analog or homolog of the EM 1 of Claim 2.

THIS PAGE BLANK (USPTO)

1/25

endo sequence from Collagen XVIII.

Sequence Range: 1-555

Nucleotide 1 = Start for Endostatin and fragments EM1 and EM2.

EM1 fragment ends at nucleotide 525, EM2 fragment ends at nucleotide 501.

```

      5      10      15      20      25      30      35      40      45
CAT ACT CAT CAG GAC TTT CAG CCA GTG CTC CAC CTG GTG GCA CTG AAC
GTA TGA GTA GTC CTG AAA GTC GGT CAC GAG GTG GAC CAC CGT GAC TTG

50      55      60      65      70      75      80      85      90      95
ACC CCC CTG TCT GGA GGC ATG CGT GGT ATC CGT GGA GCA GAT TTC CAG
TGG GGG GAC AGA CCT CCG TAC GCA CCA TAG GCA CCT CGT CTA AAG GTC

100     105     110     115     120     125     130     135     140
TGC TTC CAG CAA GCC CGA GCC GTG GGG CTG TCG GGC ACC TTC CGG GCT
ACG AAG GTC GTT CGG GCT CGG CAC CCC GAC AGC CCG TGG AAG GCC CGA

145     150     155     160     165     170     175     180     185     190
TTC CTG TCC TCT AGG CTG CAG GAT CTC TAT AGC ATC GTG CGC CGT GCT
AAG GAC AGG AGA TCC GAC GTC CTA GAG ATA TCG TAG CAC GCG GCA CGA

195     200     205     210     215     220     225     230     235     240
GAC CGG GGG TCT GTG CCC ATC GTC AAC CTG AAG GAC GAG GTG CTA TCT
CTG GCC CCC AGA CAC GGG TAG CAG TTG GAC TTC CTG CTC CAC GAT AGA

245     250     255     260     265     270     275     280     285
CCC AGC TGG GAC TCC CTG TTT TCT GGC TCC CAG GGT CAA CTG CAA CCC
GGG TCG ACC CTG AGG GAC AAA AGA CCG AGG GTC CCA GTT GAC GTT GGG

290     295     300     305     310     315     320     325     330     335
GGG GCC CGC ATC TTT TCT TTT GAC GGC AGA GAT GTC CTG AGA CAC CCA
CCC CGG GCG TAG AAA AGA AAA CTG CCG TCT CTA CAG GAC TCT GTG GGT

340     345     350     355     360     365     370     375     380
GCC TGG CCG CAG AAG AGC GTA TGG CAC GGC TCG GAC CCC AGT GGG CGG
CGG ACC GGC GTC TTC TCG CAT ACC GTG CCG AGC CTG GGG TCA CCC GCC

385     390     395     400     405     410     415     420     425     430
AGG CTG ATG GAG AGT TAC TGT GAG ACA TGG CGA ACT GAA ACT ACT GGG
TCC GAC TAC CTC TCA ATG ACA CTC TGT ACC GCT TGA CTT TGA TGA CCC

435     440     445     450     455     460     465     470     475     480
GCT ACA GGT CAG GCC TCC TCC CTG CTG TCA GGC AGG CTC CTG GAA CAG
CGA TGT CCA GTC CGG AGG AGG GAC GAC AGT CCG TCC GAG GAC CTT GTC

485     490     495     500     505     510     515     520     525
AAA GCT GCG AGC TGC CAC AAC AGC TAC ATC GTC CTG TGC ATT GAG AAT
TTT CGA CGC TCG ACG GTG TTG TCG ATG TAG CAG GAC ACG TAA CTC TTA

530     535     540     545     550     555
AGC TTC ATG ACC TCT TTC TCC AAA TAG
TCG AAG TAC TGG AGA AAG AGG TTT ATC

```

Fig. 1

THIS PAGE BLANK (USPTO)

Sequence Range: 1 to 184

5	10	15	20	25	30	35	40	45						
HTH	QDF	QPV	LHL	VAL	NTP	LSG	GMR	GIR	GAD	FQC	FQQ	ARA	VGL	SGT
50	55	60	65	70	75	80	85	90						
FRA	FLS	SRL	QDL	YSI	VRR	ADR	GSV	PIV	NLK	DEV	LSP	SWD	SLF	SGS
95	100	105	110	115	120	125	130	135						
QGQ	LQP	GAR	IFS	FDG	RDV	LRH	PAW	PQK	SVW	HGS	DPM	GRR	LME	SYC
140	145	150	155	160	165	170	175	180						
ETW	RTE	TTG	ATG	QAS	SLL	SGR	LLE	QKA	ASC	HNS	YIV	LCI	ENS	FMT

SFS K

Fig. 2

THIS PAGE BLANK (USPTO)

3/25

FIG. 3





FIG. 4



THIS PAGE BLANK (USPTO)

4/25

FIG. 5





FIG. 6



THIS PAGE BLANK (USPTO)

FIG. 7





FIG. 8




THIS PAGE BLANK (USPTO)

6/25

FIG. 9

His.endostatin (bacteria)
Endostatin (Yeast)
His.endostatin (yeast)



THIS PAGE BLANK (USPTO)

7/25




Fig. 10

THIS PAGE BLANK (USPTO)

8/25





Fig. 11

THIS PAGE BLANK (USPTO)


9/25

FIG. 12 A



Control +bFGF

FIG. 12 B



Endostatin 20 µg/ml

THIS PAGE BLANK (USPTO)

10/25





Fig. 13

THIS PAGE BLANK (USPTO)


11/25

FIG.14a




Negative control

FIG.14b



**Positive control
(VEGF)**

FIG.14c



**VEGF +
endostatin 20 µg**

THIS PAGE BLANK (USPTO)

12/25




Fig. 15A

THIS PAGE BLANK (USPTO)

13/25




Fig. 15B

THIS PAGE BLANK (USPTO)

14/25




Fig. 16

THIS PAGE BLANK (USPTO)

15/25

FIG.17_a



VEGF + endostatin (10 μ g)


FIG.17_b



**VEGF + endostatin (10 μ g) +
polyclonal antiserum (50 μ g)**

THIS PAGE BLANK (USPTO)

Fig. 18



THIS PAGE BLANK (USPTO)

17/25

FIG.19A




FIG.19B



FIG.19C



FIG.19D






FIG.19E



THIS PAGE BLANK (USPTO)



Day 1 Day 3 Day 5 Day 7 Day 9 Day 11

Fig. 20

THIS PAGE BLANK (USPTO)

19/25

Fig. 21





Fig. 22



THIS PAGE BLANK (USPTO)

20/25




Fig. 23

THIS PAGE BLANK (USPTO)

21/25

FIG.24A





FIG.24B



THIS PAGE BLANK (USPTO)

22/25

FIG.25A




FIG.25B




FIG.25C





FIG.25D



THIS PAGE BLANK (USPTO)

Construct Name	Primer Sequence	Cloning Sites	Vector	Protein Sequence
pET17b/ his.mendo	5'-GGC ATA TGC ATA CTC ATC AGG- ACT TT-3' (up) (SEQ ID NO:3)	NdeI & XhoI	Prokaryotic expression, pET	MGHHHHHHHHSSGHIDDDDKH M-mendo (SEQ ID NO:5)
	5' AAC TCG AGC TAT TTG GAG AAA- GAG GT-3' (down) (SEQ ID NO:4)			
pET28a/ mendo	5'-GGC ATA TGC ATA CTC ATC AGG- ACT TT-3' (up) (SEQ ID NO:3)	NdeI & NotI	Prokaryotic expression, pET	MGSSHHHHHHSSGLVPRGSHM- mendo (SEQ ID NO:7)
	5'-AAG CGG CCG CCT ATT TGG AGA- AAG AGG T-3' (down) (SEQ ID NO:6)			
pET28a/ EM-1	5' TTC CAT ATG CAT ACT CAT CAG- GAC TTT CAG CCA-3' (up) (SEQ ID NO:8)		Prokaryotic expression, pET	MGSSHHHHHHSSGLVPRGSHM-me ndo (SEQ ID NO:7)
	5' TTA GCG GCC GCC TAC TCA ATG- CAC AGG ACG ATG TA-3' (down) (SEQ ID NO:9)			
pET28a/ EM-2	5' TTC CAT ATG CAT ACT CAT CAG- GAC TTT CAG CCA-3' (up) (SEQ ID NO:8)		Prokaryotic expression, pET	MGSSHHHHHHSSGLVPRGSHM-me ndo (SEQ ID NO:7)
	5' TTA GCG GCC GCC TAG TTG TGG- CAG CTC GCA GCT TTC TG-3' (down) (SEQ ID NO:10)			

Fig. 26A

THIS PAGE BLANK (USPTO)

Construct Name	Primer Sequence	Cloning Sites	Vector	Protein Sequence
pPICZ α A/ mendo	5' GGG AAT TCC ATA CTC ATC AGG- ACT TT-3' (up) (SEQ ID NO:11)	<i>Eco</i> RI & <i>Not</i> I	Eukaryotic expression, yeast/pPICZ α A	EF-mendo
	5' AAG CGG CCG CCT ATT TGG AGA- AAG AGG T-3' (down) (SEQ ID NO:6)			
pPICZ α A/ His.mendo	5' AAG AAT TCC ATC ATC ATC ATC- ATC ACA GCA GC-3' (up) (SEQ ID NO:12)	<i>Eco</i> RI & <i>Not</i> I	Eukaryotic expression, yeast/pPICZ α A	EFMGHHHHHHHHSSGHIDDD KHM-mendo (SEQ ID NO:13)
	5' AAG CGG CCG CCT ATT TGG AGA- AAG AGG T-3' (down) (SEQ ID NO:6)			
pPICZ α A/ Hendo	5' TTT GAA TTC GCC CAC AGC CAC- CGC GAC TTC CAG CCG GTG CTC- CAC-3' (up) (SEQ ID NO:14)	<i>Eco</i> RI & <i>Not</i> I	Eukaryotic expression, yeast/pPICZ α A	EF-hendo
	5' AAA AGC GGC CGC CTA CTT GGA- GGC AGT CAT GAA GCT GTT CTC- AAT-3' (down) (SEQ ID NO:15)			
pPICZ α A/ Restin	5' TTT TTT GAA TTC ATT TCA AGT- GCC AAT TAT GAG AAG CCT GCT- CTG CAT- TTG-3' (up) (SEQ ID NO:16)	<i>Eco</i> RI & <i>Not</i> I	Eukaryotic (Yeast), Pichia, pPICZ α A	EF-restin
	5' AAG AAT GCG GCC GCT TAC TTC- CTA GCG TCT GTC ATG AAA CTG- TTT TCG AT-3' (down) (SEQ ID NO:17)			

Fig. 26B

THIS PAGE BLANK (USPTO)

Construct Name	Primer Sequence	Cloning Sites	Vector	Protein Sequence
pPICZ α A/ His Restin	5' AAT TCC ATC ACC ATC ACC ATC- ACG-3' (up) (SEQ ID NO:18)	<i>Eco</i> RI (oligo insertion)	Eukaryotic (Yeast), Pichia, pPICZ α A	EFHHHHHHH-restin (SEQ ID NO:20)
	5' AAT TCG TGA TGG TGA TGG TGA-TGG-3' (down) (SEQ ID NO:19)			
pET28a/ apomigren	5' TTT CAT ATG ATA TAC TCC TTT- GAT GGT CGA GAC ATA ATG ACA-3' (up) (SEQ ID NO:21)	<i>Nde</i> I & <i>Not</i> I	Prokaryotic, pET system	MGSSHHHHHHSSGLVPRGSHM-apomigren (SEQ ID NO:7)
	5' AAT GCG GCC GCT TAC TTC CTA- GCG TCT GTC ATG AAA CTG TTT- TCG AT-3' (down) (SEQ ID NO:22)			
pPICZ α A/ apomigren	5' AAG AAT TCC ATC ATC ATC ATC- ATC ACA GCA GC-3' (up) (SEQ ID NO:12)	<i>Eco</i> RI & <i>Not</i> I	Eukaryotic (Yeast), Pichia, pPICZ α A	EFMGSSHHHHHHSSGLVPRGSHM-apomigren (SEQ ID NO:23)
	5' AAT GCG GCC GCT TAC TTC CTA- GCG TCT GTC ATG AAA CTG TTT- TCG AT-3' (down) (SEQ ID NO:22)			

Fig. 26C

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/26057

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/78 C12Q1/68 C12N15/62 C07K16/18
A61K38/39

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

A	O'REILLY M S ET AL: "ENDOSTATIN: AN ENDOGENOUS INHIBITOR OF ANGIOGENESIS AND TUMOR GROWTH" CELL, vol. 88, no. 2, 24 January 1997, pages 277-285, XP002044657 cited in the application see the whole document	
---	--	--

A	L. STÄNDKER ET AL.: "Isolation and characterization of the circulating forms of human endostatin." FEBS LETTERS, vol. 420, 1997, pages 129-133, XP002100417 cited in the application see the whole document	
---	---	--

-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 April 1999

Date of mailing of the international search report

03/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hix, R

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/26057

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 17-30
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

THIS PAGE BLANK (USPTO)